



Therapeutic significance and pharmacological activities of antidiarrheal medicinal plants mention in Ayurveda: A review

Ashish Mishra, Ankit Seth, Santosh Kumar Maurya

Department of Ayurvedic Pharmacy Laboratory, Rajiv Gandhi South Campus, Banaras Hindu University, Barkachha, Mirzapur, Uttar Pradesh, India

Address for correspondence:

Santosh Kumar Maurya, Ayurvedic Pharmacy Laboratory, Rajiv Gandhi South Campus, Banaras Hindu University, Barkachha, Mirzapur - 231 001, Uttar Pradesh, India.
E-mail: dravyapharma@gmail.com

Received: February 23, 2016

Accepted: April 08, 2016

Published: May 04, 2016

ABSTRACT

Diarrhea is a serious problem affecting 3-5 billion people per year around the world, especially children of below 5 years. 70% of the world population uses traditional and indigenous medicine for their primary health care. The facts of these indigenous remedies are passed verbally and sometimes as documents. Since ancient time, Ayurveda is the main system of healing in South East Asian countries. Indian literature from ayurvedic texts and other books claim the potency of several plants in the treatment of diarrhea. As the global prospective of ayurvedic medicine is increasing, interest regarding the scientific basis of their action is parallelly increasing. Researchers are doing experiments to establish the relation between the claimed action and observed pharmacological activities. In the present article, an attempt was made to compile the scientific basis of medicinal plants used to cure diarrhea in Ayurveda. Literature was collected via electronic search (PubMed, ScienceDirect, Medline, and Google Scholar) from published articles that reports antidiarrheal activity of plants that were mentioned in Ayurveda classics. A total of 109 plant species belonging to 58 families were reported for their antidiarrheal activity. Several Indian medicinal plants have demonstrated promising antidiarrheal effects, but the studies on the antidiarrheal potentials of these plants are not taken beyond proof of concept stage. It is hoped that the article would stimulate future clinical studies because of the paucity of knowledge in this area.

KEY WORDS: Ayurveda, diarrhea, medicinal plant, traditional medicine

INTRODUCTION

Gastroenteritis is a clinico-pathological term that refers to inflammation and oxidative stress of the intestines which leads to disturbance in the balance of secretory and absorptive function of the intestines resulting in diarrhea [1,2]. Hence, diarrhea can be defined as a gastrointestinal disorder in which there is a rapid transit of gastric contents through the intestine, which is characterized by abnormal fluidity and high frequency of fecal evacuation, usually semisolid or watery fecal matter, three or more times/day [1-3]. There is an increase in flow rate of feces with or without the presence of blood and mucus, accompanied by increased secretion and decreased absorption of fluid, leading to loss of water and electrolytes [2,4]. The major causative agents of diarrhea in human beings include a variety of enteric pathogenic bacteria such as *Salmonella typhi*, *Shigella flexneri*, *Escherichia coli*, *Staphylococcus aureus*, *Vibrio cholerae*, and *Candida albicans* [4,5]. Viruses, protozoans, helminths, intestinal disorders, immunological factor, and medications can also cause diarrhea in human being [6-8]. Etiological factors for diarrhea include the food intolerances, contaminated drinking water, undercooked meat and eggs,

inadequate kitchen hygiene, poor sanitation [9], bile salts, hormones, irritable bowel syndrome, and intoxication [10]. According to the World Health Organization (WHO), diarrhea affects 3-5 billion people/year worldwide and causes 5 million deaths per annum [11]. Children, however, are more susceptible to the disease, which is the one of the leading causes of death in infants and children below 5 years of age [12].

Due to high mortality and morbidity, especially in children, the WHO together with the United Nations Children's Fund has initiated Diarrhea Disease Control Program to control diarrhea in developing countries. Oral rehydration solution [13], zinc solution [14], probiotics [15], and specific antibiotics have reduced mortality rate in diarrheal disease. However, chronic diarrhea is still a life challenging problem in some regions of the world. Unfortunately, the program does not reach to the needy, and the disease is still a major challenge in front of primary health practitioner as well as researcher. Therefore, the different traditional systems of medicines such as Chinese medicine [16], Japanese medicine [17], acupuncture therapy [18], and ayurvedic medicine [19] are included in this program.

Since ancient time's medicinal plants have been used to treat different ailments due to their accessibility, availability, inherited practice, economic feasibility, and perceived efficacy [20]. Nowadays, use of medicines from plant source increases significantly with conventional therapies. Hence, the plants are gaining more attention by the researchers to find out new and effective agents for different diseases. Several medicinal plants in the different regions of the world have been used to cure diarrhea [19,21].

The knowledge of indigenous medicines is passing from generation to generation orally worldwide [22]. It is, therefore, documentation of such knowledge as well as reported the scientific basis of their pharmacological potential is necessary since they are usually consider as free from adverse effects. A range of medicinal plants were reported for their effectiveness in diarrhea [23-27]. The protective role of these plants is probably due to their anti-inflammatory, antioxidant, and astringent properties [28]. India has a rich plant resources providing valuable medicine, which are conveniently used in Ayurveda, Unani, and other system of medicines for the treatment of various diseases [29]. Keeping this in view, the present article was initiated, with an aim to compile the scientific basis of medicinal plants used to cure diarrhea. A variety of curative agents from these indigenous plants has been isolated. These isolated compounds are belonging to different phytochemical classes such as flavonoids, saponins, terpenoids, steroids, phenolic compounds, and alkaloids [30-32]. Flavonoids and saponins inhibit the release of prostaglandins, autocoids, and contractions caused by spasmogens as well as motility and hydroelectrolytic secretions [33,34] while saponins may prevent release of histamine [35]. Polyphenols and tannins provide strength to intestinal mucosa, decrease intestinal secretion, intestinal transit and promotes balance in water transport across the mucosal cells [36].

Previously, we enumerated a large number of plants, which are used in the ayurvedic system as antidiarrheal [19]. A majority of these plants have been investigated pharmacologically with respect to the potential antidiarrheal activity. In this review, we present ethnopharmacological data of 109 plant species belonging to 58 families mentioned in ayurvedic texts for controlling diarrhea with their possible mechanism of action [Table 1 and Figure 1]. Mostly, leaf (23%), root (14%), barks (11%), fruit (9%), and seed (8%) of the plants are used for antidiarrheal activity [Figure 2].

DISCUSSION

Since ages, human beings have relied on plants as a resource of the therapeutic arsenal in the fight against certain human diseases. Plant-based drugs have formed the basis of traditional medicine systems, i.e., Ayurveda, Siddha, Unani, Homeopathy, and Chinese. Herbal-based therapy is one of the popular and effective practices to overcome the illness. The WHO also promotes utilization of local knowledge of plant-based medicines in health care. It has been reported by the WHO that about 70-80% of the population in developing countries relies

on traditional/ethno medicines/for their primary health care. Since ancient time ayurvedic system of medicine is indigenous to and widely practiced in India. Nature has bestowed India with an enormous wealth of medicinal plants. Therefore, their rational uses for combating diseases are described traditionally.

Acharya Charaka has mentioned a group of antidiarrheal plants named as *Purish-Samgarahaniya Mahakashaya*, which includes priyangu (*Callicarpa macrophylla*), ananta (*Hemidesmus indicus* R.B.), seed of amra (*Mangifera indica*), katvanga (*Ailanthus excelsa* Roxb.), lodhra (*Symplocos racemosa*), mocharasa (*Salmalia malabarica* Schott and Endl.), samanga – *Rubia cordifolia*, flower of dhataki – *Woodfordia fruticosa*, padma – lotus (*Nelumbo nucifera*), and filaments of padma – lotus (*N. nucifera*). Moreover, he also listed some most useful antidiarrheal plants such as katavanga (*A. excelsa* Roxb.), *mustaka* (*Cyperus rotundus* Linn.), *amrita* (*Tinospora cordifolia* [Willd.] Miers ex Hook. f. & Thoms.), *ativisha* (*Aconitum heterophyllum* Wall. ex. Royle.), *bilva* (*Aegle marmelos* Correa), *kumuda* (*N. nucifera* Gaertn.), *utpala*, *padma*, *kutaja* bark (*Holarrhena antidysenterica* [Linn.] Wall.), *gambhari* fruit (*Gmelina arborea* Roxb.), *prishniparni* (*Uraria picta* [Jacq.] Desv. Ex DC.), and *bala* (*Sida cordifolia*) [187]. In addition, Acharya Susuruta mentioned that the *vacha* (*Acorus calamus* Linn.) and *haridra* (*Curcuma longa* Linn.), etc., are best for *amatisara* (diarrhea where undigested food matter pass in stool) while *ambastha* (*Cissampelos pareira* Linn.) and *priyangu* (*C. macrophylla*) are best for *pakwatisara* (diarrhea where only digested food matter pass in stool) [188].

The ayurvedic Pharmacopoeia mentioned more than 1200 species of plants, nearly 100 minerals and over 100 animal products officially. Although there is no record of pharmacological testing during the period when ayurvedic texts were written. However, nowadays, extensive researches are carried out concerning the phytopharmacological basis of their therapeutic principles. Public, academic as well as government organizations are showing interest in the scientific mechanism of action exerted by these plants. Similar to modern and other traditional medicines, ayurvedic medicines have been also evaluated for their phytopharmacology with the help of advances in science and technology. Scientific screening on laboratory animal and *in vitro* evaluations supports traditional uses of medicinal plants.

In the present scenario, modern pharmaceuticals offer a number of medicines for diarrhea, but diarrhea still remains a major health threat to the people in tropical and subtropical countries. It is one of the leading causes of mortality in children especially under the age of 5 years [12]. Different factors such as infections, malnutrition, food intolerances, intestinal disorders, and some medications may trigger diarrhea [6-8]. Currently, available pharmacological treatments are seem to be insufficient in diarrhea control. It is because of lack of admittance, high cost, and adverse effects of modern pharmaceuticals as well as therapeutic approaches. Therefore, investigations on drugs from different alternative and complementary medicines along with traditional system of medicines were going on.

Table 1: Antidiarrheal medicinal plants

Sanskrit name	Botanical name	Family	Part used	Extract/dose	Standard drug and dose	Model	Mechanism	References
Atibala	<i>Abutilon indicum</i> (Linn.) Sweet.	Malvaceae	Leaf	Methanolic and aqueous extract	Loperamide (1 mg/kg)	Gastrointestinal motility test, castor oil-induced diarrhea model, and PGE ₂ -induced enteropooling model		[37]
Khadir	<i>Acacia catechu</i> Willd.	Fabaceae	Heartwood	Ethyl acetate extract (250 mg/kg)	Diphenoxylate (10 mg/kg) and atropine (1 ml/200 g, p.o.)	Castor oil-induced diarrhea model		[38]
Babool	<i>Acacia nilotica</i> Delile & Ssp. <i>indica</i> (Benth.) Brenan.	Caesalpinaceae	Bark	Petroleum ether, methanolic and aqueous extract		Castor oil- and magnesium sulfate-induced diarrhea and barium chloride-induced gastrointestinal motility test		[39]
Ativisha	<i>Aconitum heterophyllum</i> Wall. ex. Royle.	Ranunculaceae	Root	Ethanol extract (50, 100, and 200 mg/kg) and isolated aconitine	Loperamide (2 mg/kg, p.o.) and atropine (0.1 mg/kg, s.c.)	Castor oil-induced diarrhea model, Small intestinal transit time, PGE ₂ -induced enteropooling, and gastric emptying test	Prevented Na ⁺ and K ⁺ loss	[40]
Vacha	<i>Acorus calamus</i> Linn.	Araceae	Root and essential oil	Methanolic extract and n-hexane fraction		Castor oil-induced diarrhea model, spasmolytic activity	Inhibition of spontaneous and High K ⁺ -induced contractions and antispasmodic action	[41,42]
Bilva	<i>Aegle marmelos</i> Correa.	Rutaceae	Rhizome Unripe fruit pulp	Aqueous and methanolic extract (3, 7.5, and 15 mg) Aqueous extract	-	Castor oil-induced diarrhea model Antimicrobial activity	Through reduced bacterial adherence to intestinal wall and Invasion of Hep-2 cells	[43-50]
			Leaf	Aqueous extract (50, 100, and 200 mg/kg)	Loperamide (3 mg/kg orally)	Castor oil-induced diarrhea, magnesium sulfate-induced diarrhea, and gastric transit time		
			Fruit	Polyherbal formulation (25, 50, and 100 mg/kg)	Mebarid (10 ml/kg, p.o.)	Castor oil-induced diarrhea model, intestinal secretion model, and antispasmodic effect	Inhibition of intestinal transit of food material and inhibition of intestinal secretion	
			Unripe fruit	Aqueous and methanolic extract		Castor oil-induced diarrhea model		
			Fruit	Aqueous extract	Diphenoxylate and yohimbine	Castor oil-induced diarrhea model		
			Dried fruit pulp	Ethanol extract	-	<i>In vitro</i> antibacterial activity		
			Root	Chloroform extract	-	Castor oil-induced diarrhea model		

Contd...

Table 1: (Contd....)

Sanskrit name	Botanical name	Family	Part used	Extract/dose	Standard drug and dose	Model	Mechanism	References
Gorakghanja	<i>Aerva lanata</i>	Amaranthaceae	Unripe fruit	(50, 100 mg/kg)	-	Gastrointestinal motility test and castor oil-induced diarrhea model		[51]
Aralu	<i>Ailanthus excelsa</i> Roxb.	Simaroubaceae	Whole plant	Alcoholic and aqueous extract (400 and 800 mg/kg)	Loperamide (3 mg/kg, i.p.)	Castor oil-induced diarrhea model		[52]
Shirish	<i>Albizia lebeck</i> Benth.	Caesalpinaceae	Bark	Chloroform, aqueous and ethanolic extract	Atropine (0.1 mg/kg, i.p.)	Castor oil-induced diarrhea model		[53]
Saptaparna	<i>Alstonia scholaris</i> R.Br.	Apocynaceae	Seed	Crude extract	Loperamide (1 mg/kg, i.p.)	Castor oil-induced diarrhea model		[54,55]
			Bark	Methanolic extract	Loperamide	Castor oil-induced diarrhea model	Spasmodic activity mediated possibly through CCB	
Tanduliya	<i>Amaranthus spinosus</i> Linn.	Amaranthaceae	Whole plant	Ethanollic extract (250 mg/kg)	Loperamide (50 mg/kg)	Castor oil-induced diarrhea model		[56]
Eshwari	<i>Aristolochia indica</i> Linn.	Aristolochiaceae	Root	Ethanollic extract (200, and 400 mg/kg)	Yohimbine	Castor oil-induced diarrhea model, charcoal-induced gastrointestinal motility test		[57]
Shatavari	<i>Asparagus racemosus</i> Willd.	Liliaceae	Root	Aqueous extract and ethanolic extract	Diphenoxylate (5 mg/kg, p.o.)	Castor oil-induced diarrhea model, charcoal-induced gastrointestinal motility test		[58]
Hijjala	<i>Barringtonia acutangula</i> (Linn.) Gaertn.	Lecythidaceae	Root	Ethanollic extract (200 and 400 mg/kg)	Loperamide (3 mg/kg, p.o.)	Castor oil-induced diarrhea model, gastrointestinal tract motility, PGE ₂ -induced enteropooling		[59,60]
Kovidara	<i>Bauhinia purpurea</i> Linn.	Caesalpinaceae	Leaf and seed	Methanollic extract (200 and 400 mg/kg)	Diphenoxylate (5 ml/kg, p.o.)	Castor oil-induced diarrhea model		[61]
Kanchnar	<i>Bauhinia variegata</i> Linn.	Caesalpinaceae	Leaf	Ethanollic extract	Loperamide (3 mg/kg, p.o.)	Castor oil and magnesium sulfate-induced diarrhea models		[62]
Daruharidra	<i>Berberis aristata</i> DC.	Berberaceae	Stem bark	(250, 500, and 1000 mg/kg, p.o.)	Loperamide (2 mg/kg, po)	Castor oil-induced diarrhea model	Inhibit the intestinal secretory response	[63-65]
			Stem	Aqueous extract	Loperamide (25 mg/kg)	Castor oil-induced diarrhea model		
			Bark	Ethanollic, aqueous extract and isolated berberine	Loperamide (3 mg/kg orally)	Magnesium sulfate-induced diarrhea, castor oil-induced intestinal secretions		[66]
Sinduri	<i>Bixa orellana</i> Linn.	Bixaceae	Stem	Ethyl alcohol extract (250, 500 mg/kg)	Loperamide (3 mg/kg orally)	Castor oil-induced diarrhea model		
			Leaf	Methanollic extract (125, 250, and 500 mg/kg)	Loperamide (3 mg/kg orally)	Castor oil-induced diarrhea model		

Contd...

Table 1: (Contd....)

Sanskrit name	Botanical name	Family	Part used	Extract/dose	Standard drug and dose	Model	Mechanism	References
Sallaki	<i>Boswellia serrata</i> Roxb. Ex Coleb.	Burseraceae	Gum resin	Hydroalcoholic extract and 3-acetyl-11-keto- β -boswellic acid	Atropine (1 mg/kg)	Upper gastrointestinal transit in croton oil-treated animal, castor oil-induced diarrhea model	Inhibition of acetylcholine-induced contractions by the L-type Ca^{2+} -channel blockers	[67]
Parnabija	<i>Bryophyllum pinnatum</i> (Lam.) Kurz.	Crassulaceae	Leaf	Aqueous extract (100, 200, and 300 mg/kg)	Loperamide (5 mg/kg)	Castor oil-induced diarrhea model, castor oil-induced enteropooling, small intestinal transit time		[68]
Priyala	<i>Buchanania lanzan</i> Spreng.	Fagaceae	Leaf	(200 and 400 mg/kg)	Loperamide (1 mg/kg)	Castor oil-induced diarrhea model, charcoal meal test	Inhibition Na^+K^+ ATPase activity	[69]
Palash	<i>Butea monosperma</i> Lam. Kuntze.	Fabaceae	Stem bark	Ethanollic extract	Loperamide (3 mg/kg orally)	Castor oil-induced diarrhea model and PGE ₂ -induced enteropooling		[70]
Latakaranja	<i>Caesalpinia bonducella</i> Flem.	Caesalpinaceae	Leaf	Methanolic extract and its ethyl acetate, chloroform, and petroleum ether fractions (200 and 400 mg/kg)	Loperamide (5 mg/kg, p.o.)	Castor oil-induced diarrhea model	Antibacterial activity	[71]
Gumohar	<i>Caesalpinia pulcherrima</i> L.	Caesalpinaceae	Bark	Ethanollic extract (500 mg/kg)	Loperamide (50 mg/kg orally)	Castor oil-induced diarrhea model		[72]
Arka	<i>Calotropis gigantea</i> R.Br.	Asclepiadaceae	Aerial part	Hydroalcoholic extract (200 and 400 mg/kg)	Atropine (3 mg/kg, i.p.)	Castor oil-induced diarrhea model		[73]
Arka	<i>Calotropis procera</i> (Ait.) R.Br.	Asclepiadaceae	Dry latex	(500 mg/kg)	Atropine (0.1 mg/kg, i.p.)	Castor oil-induced enteropooling, electrolyte concentration in the intestinal fluid and intestinal transit		[74-77]
Tea	<i>Camellia sinensis</i> (Linn.) O. Kuntze.	Theaceae	Leaf	Ethanollic extract (250, 500 mg/kg)	Loperamide (4 mg/kg)	Castor oil-induced diarrhea model		
Hinsra	<i>Capparis zeylanica</i> Linn.	Capparidaceae	Leaf	Methanollic extract	Loperamide (3 mg/kg orally)	Castor oil-induced diarrhea model		
Erand karkati	<i>Carica papaya</i> Linn.	Caricaceae	Fruit	Aqueous and alcoholic extract (100, 200 mg/kg)		Castor oil-induced diarrhea model, charcoal meal test, enteropooling method		
Shitiwar	<i>Celosia argentea</i> Linn.	Amaranthaceae	Leaf	Aqueous extract	Castor oil-induced diarrhea model and PGE ₂ -induced enteropooling			[78]
Patra	<i>Cinnamomum tamala</i> Buch.-Ham.	Lauraceae	Bark	Methanollic extract (100, 200 mg/kg)	Loperamide (3 mg/kg orally)	Castor oil-induced diarrhea model and small intestine transit method		[79]
Twaka	<i>Cinnamomum zeylanicum</i> Linn.	Lauraceae	Bark	Alcoholic and aqueous extract (100, 200 and 400 mg/kg)	Loperamide (3 mg/kg, p.o.)	Castor oil-induced diarrhea model and magnesium sulfate-induced diarrhea		[80]
				Alcoholic extract (100, 200 mg/kg)	Atropine (0.1 mg/kg, s.c)	Castor oil-induced diarrhea model, charcoal meal test, PGE ₂ -induced diarrhea		[81]
				Ethanollic extract (25, 50, and 100 mg/kg)	Loperamide (5 mg/kg)	Castor oil-induced diarrhea model and magnesium sulfate-induced diarrhea		[82]
				Aqueous extract (100 and 200 mg/kg)	Loperamide (5 mg/kg)	Castor oil-induced diarrhea model and magnesium sulfate-induced diarrhea		[83]

Contd...

Table 1: (Contd....)

Sanskrit name	Botanical name	Family	Part used	Extract/dose	Standard drug and dose	Model	Mechanism	References
Patha	<i>Cissampelos pareira</i> Linn.	Menispermaceae	Root	Ethanollic extract (25-100 mg/kg)		Castor oil-induced diarrhea model	Inhibitory effect on the concentration of Na ⁺ and K ⁺ , reduction in the lipid peroxidation and Prevention from oxidative stress	[84]
Hulhul	<i>Cleome viscosa</i> L.	Capparidaceae	Whole plant	Methanolic extract	Diphenoxylate (5 mg/kg orally)	Castor oil-induced diarrhea model and PGE ₂ -induced enteropooling gastrointestinal motility		[85]
Vaamana-haati	<i>Clerodendrum indicum</i>	Verbenaceae	Leaf	Methanolic extract and chloroform fraction	-	Castor oil-induced diarrhea model		[86]
Aparajita	<i>Clitoria ternatea</i> L.	Fabaceae	Leaf	Methanolic extract (100, 200, and 300 mg/kg)	Loperamide (3 mg/kg)	Castor oil-induced diarrhea model and small intestine transit method		[87,88]
			Root	Alcoholic extract (100, 200, and 400 mg/kg)	Atropine (5 mg/kg, i.p.)	Castor oil-induced diarrhea model, intestinal transit and castor oil-induced enteropooling		
Dhanyaka	<i>Coriandrum sativum</i> Linn.	Apiaceae	Leaf	Aqueous extract (150 and 300 mg/kg)	Loperamide (3 mg/kg)	Castor oil-induced diarrhea model		[89]
Varuna	<i>Crataeva nurvala</i> Buch.-Ham.	Capparidaceae	Stem bark	Ethanollic extract (500 mg/kg)		Castor oil-induced diarrhea model, castor oil-induced enteropooling, and small intestine transit model		[90]
Jiraka	<i>Cuminum cyminum</i> Linn.	Apiaceae	Seed	Aqueous extract	Loperamide (3 mg/kg)	Castor oil induce diarrhea model, PGE ₂ -induced enteropooling model, intestinal transit by charcoal		[91]
Haridra	<i>Curcuma longa</i> Linn.	Zingiberaceae	Rhizome	Aqueous extract (200 mg/kg)		Castor oil-induced diarrhea model		[92]
Durva	<i>Cynodon dactylon</i> Pers.	Poaceae	Whole plant	Methanolic extract (200 and 300 mg/kg)	Atropine (5 mg/kg orally)	Castor oil-induced diarrhea model, gastrointestinal charcoal meal test, and enteropooling model		[93]
Mustaka	<i>Cyperus rotundus</i> Linn.	Cyperaceae	Rhizome	Methanolic extract (250-500 mg/kg) Aqueous extract	-	Castor oil-induced diarrhea model		[94,95]
Goraksha	<i>Dalbergia lanceolaria</i> Linn.f.	Fabaceae	Bark	Petroleum ether, ethanollic extract	Diphenoxylat (5 mg/kg, p.o.)	Antibacterial activity against EPEC and EIEC and Shigella flexneri	Antibacterial, antiangiardial and antiprotaviral activities	
Shimsapa	<i>Dalbergia sissoo</i> Roxb. ex DC.	Fabaceae	Leaf	Ethanollic extract		Castor oil and magnesium sulfate-induced diarrhea		[96]
Kusha	<i>Desmostachya bipinnata</i> L.	Poaceae		Alcohol aqueous extract (200, 400 mg/kg)	Loperamide (3 mg/kg, p.o.)	Castor oil-induced diarrhea model and magnesium sulfate model		[97]
Virataru	<i>Dichrostachys cinerea</i> W. & A.	Mimosaceae	Leaf bark and root	Ethanollic extract (200 and 400 mg/kg)		MgSO ₄ -induced diarrhea		[98]
Tinduka	<i>Diospyros peregrina</i> Gruke.	Ebenaceae	Bark and seed	Ethanollic extract (250 and 500 mg/kg)	Loperamide (5 mg/kg, p.o.)	Castor oil-induced diarrhea model, gastrointestinal motility test with charcoal meal test		[99]
						Castor oil-induced model and small intestinal transit model		[100]
						Castor oil-induced diarrhea model		

Contd...

Table 1: (Contd....)

Sanskrit name	Botanical name	Family	Part used	Extract/dose	Standard drug and dose	Model	Mechanism	References
Amalaki	<i>Emblica officinalis</i> Gaertn.	Euphorbiaceae	Fruit	Ethanollic extract (500 mg/kg)	Loperamide (3 mg/kg)	Castor oil-induced diarrhea model	Inhibition of intestinal motility, antimicrobial action, and antisecretory effects	[101-103]
Paribhadra	<i>Erythrina indica</i> Lam.	Fabaceae	Leaf	Crude extract (500-700 mg/kg) Methanolic extract	Loperamide (10 mg/kg)	Castor oil-induced diarrhea model and enteropooling model	Mediated possibly through dual blockade of muscarinic receptors and Ca ²⁺ channels	[104]
Dugdihika Big	<i>Euphorbia hirta</i> Linn.	Euphorbiaceae	Whole plant	Ethanollic and aqueous extract (500 mg/kg)	Loperamide (5 mg/kg)	Diarrhea-induced by castor oil and magnesium sulfate, gastrointestinal motility in charcoal meal tests, and PGE ₂ -induced enteropooling		[105]
Kapittha	<i>Feronia limonia</i> Linn. Swingle	Rutaceae	Leaf	Ethanollic extract (250, 500 mg/kg) Ethanollic extract (500 mg/kg) Methanollic (3, 7.5 and 15 mg/kg)	Loperamide (50 mg/kg) Loperamide (25 mg/kg)	Castor oil-induced diarrhea model, intestinal tract motility model Castor oil-induced diarrhea model, PGE ₂ -induced enteropooling, gastrointestinal motility in both BaSO ₄ and charcoal meal tests		[106,107]
Vata	<i>Ficus benghalensis</i>	Moraceae	Leaf	Ethanollic extract (400 mg/kg)	Diphenoxylate (5 mg/kg, p.o.)	Castor oil-induced diarrhea model	Increasing colonic water and electrolyte re-absorption or by inhibiting intestinal motility	[108,109]
Kakodumbara	<i>Ficus hispida</i> Linn.	Moraceae	Leaf	Methanollic extract	Diphenoxylate (5 mg/kg, p.o.)	Gastrointestinal motility in charcoal meal test, castor oil-induced diarrhea model, and PGE ₂ -induced enteropooling		[110]
Udumbara	<i>Ficus racemosa</i> Linn.	Moraceae	Bark	Ethanollic extract 400 mg/kg	Diphenoxylate (5 mg/kg, p.o.)	Castor oil-induced enteropooling model and PGE ₂ -induced enteropooling model		[109]
Ashvattha	<i>Ficus religiosa</i> Linn.	Moraceae	Stem bark	Hydroalcoholic, acetone extract	Loperamide (3 mg/kg, p.o.)	Castor oil-induced diarrhea model		[111]
Udumber	<i>Ficus glomerata</i> L.	Moraceae	Leaf	Methanollic extract (100 and 200 mg/kg)	Atropine (3 mg/kg)	Castor oil-induced diarrhea model, castor oil-induced enteropooling, and intestinal transit		[112]
Parpata	<i>Fumaria parviflora</i>	Papeveraceae	Aerial part	Aqueous and methanollic extract	Dicyclomine, (50 and 100 mg/kg) and loperamide (10 mg/kg, p.o.)	Castor oil-induced diarrhea model	CCB blockade of muscarinic receptors	[113]

Contd...

Table 1: (Contd....)

Sanskrit name	Botanical name	Family	Part used	Extract/dose	Standard drug and dose	Model	Mechanism	References
Kasmari	<i>Gmelina arborea</i> Roxb.	Verbenaceae	Root	Ethanollic and N-butanol (200, 400 mg/kg) Aqueous and methanollic extract (0.5, 1.0 mg/ml) Methanollic extract (500-1500 mg/kg)	Loperamide (3 mg/kg, p.o.) Loperamide (5 mg/kg)	Castor oil-induced diarrhea model Castor oil-induced diarrhea model Castor oil-induced diarrhea model	Inhibition of intestinal motility and bactericidal activity	[114,115]
Sariva	<i>Hemidesmus indicus</i> R.Br.	Apocynaceae	Root	Aqueous and ethanollic extract (100 and 200 mg/kg) Ethanollic extract (200-800 mg/kg), Isolated alkaloid Ethanollic extract (250 and 500 mg/kg) Aqueous extract (400 mg/kg)	Loperamide (3 mg/kg)	Charcoal meal test and enteropooling model		
Kutaja	<i>Holarrhena antidysenterica</i> (Linn.) Wall.	Asclepiadaceae	Seed			Castor oil-induced diarrhea model, EPEC <i>in vitro</i>		[118]
Cirabilva	<i>Holoptelea integrifolia</i> Planch.	Urticaceae	Leaf		Loperamide (3 mg/kg, p.o.)	Castor oil and magnesium sulfate-induced diarrhea model		[119]
Bandhuka	<i>Ixora coccinea</i> Linn.	Rubiaceae	Flower		Loperamide (5 mg/kg)	Castor oil-induced diarrhea model		[120,121]
Vyaghra errand	<i>Jatropha curcas</i> Linn.	Euphorbiaceae	Root	Aqueous extract (400 mg/kg) Methanollic extract (50 and 100 mg/kg)	Loperamide (5 mg/kg) Chlorpromazine (30 mg/kg, i.p.)	Castor oil-induced diarrhea model Castor oil or magnesium sulfate-induced diarrhea	Inhibition of prostaglandin biosynthesis and reduction of osmotic pressure, decreases in peristaltic activity, Castor oil-induced permeability changes in intestinal mucosal membrane to water and electrolyte	[122,123]
Madhuca	<i>Madhuca indica</i> J. F. Gmel.	Sapotaceae	Dried bark	Petroleum ether and methanollic extract Ethanollic extract (250 and 500 mg/kg) Methanollic extract	Chlorpromazine (30 mg/kg, i.p.) Loperamide (50 mg/kg)	Castor oil-induced diarrhea model, gastrointestinal motility after charcoal meal Castor oil-induced diarrhea model	Inhibition of prostaglandin biosynthesis and reduction of osmotic pressure, decreases in peristaltic activity, Castor oil-induced permeability changes in intestinal mucosal membrane to water and electrolyte Inhibition of prostaglandin biosynthesis and reduction of propulsive movement of small intestine	[124]
Amra	<i>Mangifera indica</i> Linn.	Anacardiaceae	Stem bark and root bark	Methanollic extract (3, 7.5, and 15 mg/kg)		Castor oil-induced diarrhea model	By increasing colonic water and electrolyte reabsorption or by inhibiting intestinal motility	[108,125-127]
			Seed	Alcoholic and aqueous extract	Loperamide	Castor oil-induced diarrhea model		
			Leaf	Aqueous extract (25 and 50 mg/kg)	Loperamide (2 mg/kg)	Castor oil-induced diarrhea model	Enhancement of Na ⁺ -K ⁺ ATPase activity	
			Seed	Methanollic and aqueous extract (250 mg/kg)	Loperamide (3 mg/kg, p.o.)	Castor oil- and magnesium sulfate-induced diarrhea model		

Contd...

Table 1: (Contd....)

Sanskrit name	Botanical name	Family	Part used	Extract/dose	Standard drug and dose	Model	Mechanism	References
Pudina	<i>Mentha longifolia</i> (Linn.) Huds.	Lamiaceae	Whole plant	Crude extract, petroleum spirit fraction, aqueous fraction (100-1000 mg/kg)	Loperamide	Castor oil-induced diarrhea model	Inhibition of spontaneous and high K ⁺ -induced contractions, spasmodytic activity, mediated possibly through CCB	[128,129]
Lajjalu	<i>Mimosa pudica</i> Linn.	Mimosaceae	Leaf	Essential oil (20-80 mg/kg) Ethanol extract (200 and 400 mg/kg)	Loperamide (3 mg/kg, p.o.)	Castor oil-induced diarrhea model and PGE ₂ -induced enteropooling, gastrointestinal motility in charcoal meal test		[130,131]
Karvellaka	<i>Momordica charantia</i> Linn.	Cucurbitaceae	Leaf	Ethanol and aqueous extract (150 and 250 mg/kg) Aqueous extract		Gastrointestinal motility in charcoal meal test		[132]
Shobhanjana	<i>Moringa oleifera</i> Lam.	Moringaceae	Leaf	Hydroalcoholic extract (2500 mg/kg)	Loperamide (3 mg/kg)	Castor oil-induced diarrhea model, gastrointestinal transit, intestinal fluid accumulation and gastric emptying		[133,134]
Surabhi-nimba	<i>Murraya koenigii</i> (Linn.) Spreng.	Rutaceae	Leaf	Ethanol extract (150 and 300 mg/kg) Aqueous extract (200 mg/kg) and alcoholic extract (400 mg/kg)	Loperamide (3 mg/kg, p.o.) Loperamide (2 mg/kg)	Castor oil-induced diarrhea model, charcoal meal test, and PGE ₂ -induced diarrhea		[135,136]
Kamini	<i>Murraya paniculata</i> (L.) Jack.	Rutaceae	Leaf	Ethanol extract (300 and 600 mg/kg)	Loperamide (50 mg/kg)	Castor oil-induced diarrhea model		[137]
Kadali	<i>Musa paradisiaca</i> Linn.	Musaceae	Sap	0.25, 0.50, and 1.00 mL	Loperamide (2.5 mg/kg) Atropine (2.5 mg/kg)	Castor oil-induced diarrhea model, castor oil-induced enteropooling, and gastrointestinal motility		[138]
Jatiphala	<i>Myristica fragrans</i> Houtt.	Myristicaceae	Flower bud	Aqueous extract and petroleum ether extract	Atropine (2.5 mg/kg)	Antispasmodic	Inhibited the contraction produced by acetylcholine, Histamine, and prostaglandin	[139]
Kamala	<i>Nelumbo nucifera</i> Gaertn.	Nymphaeaceae	Rhizome	(100, 200, 400, and 600 mg/kg)		Castor oil-induced diarrhea model and PGE ₂ -induced enteropooling and charcoal meal test		[140-142]

Contd...

Table 1: (Contd....)

Sanskrit name	Botanical name	Family	Part used	Extract/dose	Standard drug and dose	Model	Mechanism	References
Trivrita	<i>Operculina tupeethum</i> . (Linn.) Silva Manso.	Convolvulaceae	Root bark Rhizome	Methanolic extract (100, 200, 400, and 600 mg/kg) Crude hexane extract	Atropine (0.1 mg/kg) Diphenoxylate (5 mg/kg)	Castor oil-induced diarrhea and PGE ₂ -induced enteropooling <i>In vitro</i> antibacterial activity against <i>Staphylococcus aureus</i> , <i>Pseudomonas aeruginosa</i> , <i>Salmonella typhimurium</i> , <i>Escherichia coli</i> , and <i>Enterobacter aerogenes</i>	Possibly through the presence of Ca ⁺⁺ antagonist	[143]
Syonaka	<i>Oroxylum indicum</i> Vent.	Bignoniaceae	Stem bark	Methanolic extract (400 mg/kg)	Loperamide (10 mg/kg, p.o.) Loperamide (66.67 µg/kg, p.o.)	Castor oil-induced diarrhea model Castor oil-induced diarrhea model		[144,145]
Cangeri	<i>Oxalis corniculata</i> Linn.	Gerniaceae	Bark	Flavonoids rich fraction	Loperamide (10 mg/kg, p.o.) Loperamide (66.67 µg/kg, p.o.)	Castor oil and magnesium sulfate-induced diarrheal models, barium chloride, and acetylcholine-induced intestinal contraction	Alteration of intestinal motility through modification in L- type Ca ²⁺ - channels	[146]
Gandhaprasharni	<i>Paederia foetida</i> Linn.	Rubiaceae	Root	Aqueous and Methanolic extract (160, 320, and 640 mg/kg) Ethanol extract (100, 250, and 500 mg/kg)	Atropine (5 mg/kg)	Castor oil-induced diarrhea model, magnesium sulfate-induced diarrhea, gastrointestinal motility with barium sulfate milk, cisplatin-induced gastrointestinal motility, morphine-induced reduction of motility		[147]
Pind kharjura	<i>Phoenix dactylifera</i> Linn.	Palmaaceae	Fruit	Aqueous extract (1000 and 1500 mg/kg)	Loperamide (5 mg/kg)	Castor oil-induced diarrhea model, enteropooling model, and gastrointestinal motility test		[148]
Maricha	<i>Piper nigrum</i> L.	Piperaceae	Fruit Fruit	Piperine Aqueous extract (75, 150, and 300 mg/kg) Aqueous extract (300 mg/kg)	Loperamide (2 mg/kg, p.o.) Atropine (5 mg/kg, i.p.) Chlorpromazine (30 mg/kg, i.p.) Isosorbide dinitrate (150 mg/kg, p.o.) Glibenclimide (1 mg/kg, p.o.) Yohimbine (1 mg/kg, s.c.)	Castor oil and magnesium sulfate-induced diarrhea charcoal meal test and castor oil-induced intestinal secretions Castor oil-induced diarrhea model	On α ₂ adrenergic receptors, potassium channels, and nitric oxide pathway	[149-153]

Contd...

Table 1: (Contd....)

Sanskrit name	Botanical name	Family	Part used	Extract/dose	Standard drug and dose	Model	Mechanism	References
Karkatasringi	<i>Pistacia integerrima</i> (J. L. Stewart ex Brandis)	Anacardiaceae	Gall	Piperine (8 and 32 mg/kg) Piperine (10 mg/kg)	Loperamide (10 mg/kg)	Castor oil, MgSO ₄ and arachidonic acid-induced diarrhea, castor oil induced enteropooling Castor oil-induced diarrhea model	Inhibitory effect on prostaglandins Concentration-dependent inhibition of spontaneous contractions, CCB effect. Piperine (10-100 μm) caused a rightward shift in the Ca ⁺⁺ concentration-response curves in Ca ⁺⁺ -free medium	[154]
Karanja	<i>Pongamia pinnata</i> (Linn.) Pierre.	Fabaceae	Leaf	Methanolic extract (700 and 900 mg/kg) Aqueous extract	-	Castor oil-induced diarrhea model, isolated rabbit jejunum Antibacterial, anti-giardial and antitroviral activity	Inhibits adherence of EPEC and invasion of EIEC and <i>Shigella flexneri</i> to epithelial cells	[155]
Peruka	<i>Psidium guajava</i> Linn.	Myrtaceae	Leaf	Aqueous extract	Loperamide (10 mg/kg, p.o.)	Castor oil-induced diarrhea model		[156-158]
Bijaka	<i>Pterocarpus marsupium</i>	Fabaceae	Heartwood	Aqueous extract (50-400 mg/kg) Methanolic and aqueous extract (100 mg/kg)	Loperamide (1 mg/kg, i.p.) Loperamide (5 mg/kg, p.o.)	Gastrointestinal Motility, castor oil-induced diarrhea model, and PGE ₂ -induced enteropooling Castor oil and charcoal-induced gastrointestinal motility test, intestinal transit of charcoal meal		[159]
Dadima	<i>Punica granatum</i> Linn.	Punicaceae	Seed	Ethanol extract (250 and 500 mg/kg) Methanolic extract	Mebarid (10 ml/kg, po)	Castor oil-induced diarrhea and PGE ₂ -induced enteropooling Castor oil-induced diarrhea model, spontaneous movement of the isolated rat ileum, acetylcholine-induced contractions test Castor oil-induced diarrhea model, intestinal secretion, and charcoal meal test	Antimotility and antisecretory activity	[160-162]
Mayaphala	<i>Quercus infectoria</i>	Fagaceae	Peels	Aqueous extract (100, 200, 300, and 400 mg/kg)	Loperamide (3 mg/kg, p.o.)	Castor oil-induced diarrhea model, intestinal secretion, and charcoal meal test		[163]
Sarpagandha	<i>Rauwolfia serpentina</i> Benth. ex Kurz.	Apocynaceae	Rinds of fruit	Polyherbal formulation Aqueous extract	Methanolic extract (100, 200, and 400 mg/kg) Ethanol extract (50, 100 mg/kg)	Castor oil-induced diarrhea model, intestinal secretion, and charcoal meal test		[164]
Manjistha	<i>Rubia cordifolia</i> L.	Rubiaceae	Root	Ethanol extract (50, 100 mg/kg)	Loperamide (5 mg/kg, p.o.)	Castor oil-induced diarrhea model, gastrointestinal transit time	Decrease in both sodium and potassium excretion in the intestine	[165]

Contd...

Table 1: (Contd....)

Sanskrit name	Botanical name	Family	Part used	Extract/dose	Standard drug and dose	Model	Mechanism	References
Chandan	<i>Santalum album</i> Linn.	Santalaceae	Heartwood	Methanolic extract (200, 400, and 800 mg/kg)		Castor oil-induced diarrhea model	Spasmolytic role relaxed the acetylcholine-induced, 5-HT-induced and K ⁺ -induced contractions	[166]
Asoka	<i>Saraca asoca</i> (Roxb.) De Wilde	Caesalpinaceae	Stem bark	Hydroalcoholic, acetone extract (200 mg/kg)	Loperamide (3 mg/kg, p.o.)	Castor oil-induced diarrhea model		[167]
Kushtha	<i>Saussurea lappa</i> Clarke	Asteraceae	Essential oil	100, 300, and 500 mg/kg	Loperamide (5 mg/kg)			[168]
Raj Bala	<i>Sida rhombifolia</i>	Malvaceae	Root	Methanolic extract (200 and 400 mg/kg)	Diphenoxylate (5 mg/kg)	Castor oil-induced diarrhea model, intestinal transit, and castor oil-induced intestinal fluid accumulation (enteropooling)		[169]
Kupilu	<i>Strychnos nux-vomica</i> Linn. f.	Loganiaceae	Root bark	Aqueous and Methanolic extract (3, 7.5, and 15 mg)		Castor oil-induced diarrhea model		[42]
Kataka	<i>Strychnos potatorum</i> Linn.	Loganiaceae	Seed	Methanolic extract	Diphenoxylate (5 mg/kg)	Castor oil-induced diarrhea model, effects on gastrointestinal motility and PGE ₂ -induced gastric enteropooling		[170]
Lodhra	<i>Symplocos racemosa</i> Roxb.	Symplocaceae	Bark	Ethylacetate chloroform, n-butanol and aqueous fraction (300, 500 mg/kg)		Spontaneous movement of the isolated rabbit intestine		[171]
Jambu	<i>Syzygium cumini</i> Linn. Skeels	Myrtaceae	Seed	Aqueous extract (125, 250, and 500 mg/kg)	Loperamide (2 mg/kg, p.o.)	Castor oil-induced diarrhea model, charcoal meal test, castor oil-induced intestinal secretions		[172]
Sharpunkha	<i>Tephrosia purpurea</i> (Linn.) Pers.	Fabaceae	Whole plant	Methanolic extract (300 mg/kg)	Verapamil (50 mg/kg)	Castor oil-induced diarrhea model		[173]
Arjuna	<i>Terminalia arjuna</i> (Roxb.) W. & A.	Combretaceae	Bark	Methanolic extract (100, 200, and 400 mg/kg)	Loperamide (3 mg/kg)	Castor oil and gastro intestinal motility test		[174]
Bibhitaki	<i>Terminalia bellirica</i> Roxb.	Combretaceae	Fruit	Aqueous and ethanolic extract (143, 200, and 334 mg/kg)	Loperamide (3 mg/kg)	Castor oil-induced diarrhea model, PGE ₂ -induced enteropooling and gastrointestinal motility test		[175]
Parisha	<i>Thespesia populnea</i> Soland. Ex. Correa	Malvaceae	Stem bark	Methanolic fraction (100 mg/kg) and residue fraction (10, 25, and 50 mg/kg) of aqueous extract	Loperamide (3 mg/kg)	Castor oil-induced diarrhea model, PGE ₂ -induced enteropooling, charcoal meal test	Inhibition of elevated prostaglandin biosynthesis, reduced propulsive movement of the intestine	[176,177]
				Aqueous extract (100, 200, and 400 mg/kg) and alcoholic extract (50, 100, and 200 mg/kg)	Atropine (3 mg/kg)	Castor oil-induced diarrhea model; PGE ₂ -induced enteropooling, charcoal meal test		

Contd...

Table 1: (Contd....)

Sanskrit name	Botanical name	Family	Part used	Extract/dose	Standard drug and dose	Model	Mechanism	References
Guduchi	<i>Tinospora cordifolia</i> (Willd.) Miers ex Hook.f. & Thoms.	Menispermaceae	Stern	Ethanollic and aqueous extract	Loperamide (3 mg/kg, p.o.)	Castor oil and magnesium sulfate-induced diarrhea		[178]
Adhapushpi	<i>Trichodesma indicum</i> R.Br.	Boraginaceae	Root	Ethanollic extract		Castor oil-induced diarrhea model charcoal meal transit time, castor oil-induced enteropooling		[179]
Methika	<i>Trigonella foenum-graecum</i> Linn.	Fabaceae	Whole plant	Aqueous extract (100, 200 mg/kg)	Loperamide (1 mg/kg, i.p.)	Castor oil-induced diarrhea model		[180]
Pind tagar	<i>Valeriana harlowickii</i> Wall.	Valerianaceae	Rhizome	Aqueous-Methanollic extract	Loperamide (10 mg/kg)	Castor oil-induced diarrhea model	Inhibited K ⁺ -induced contractions (0.01-0.3 mg/ml), CCB	[181]
Sampushpa	<i>Vinca major</i> L.	Apocynaceae	Aerial part	Ethanollic extract (250, 500, and 1000 mg/kg)	Loperamide (3 mg/kg, p.o.) Atropine (5 mg/kg, i.p.)	Castor oil-induced diarrhea model, castor oil and magnesium sulfate-induced enteropooling, gastrointestinal motility test using charcoal meal methods		[182]
Kutaja	<i>Wrightia tinctoria</i> Roxb. R.Br.	Apocynaceae	Bark	Ethanollic extract (500 and 1000- 189 mg/kg) and isolated steroidal alkaloid fraction (50 and 100 mg/kg)	Loperamide (0.5 mg/kg), atropine (0.1 mg/kg, i.p.)	Castor oil-induced diarrhea model, charcoal meal, PGE ₂ -induced enteropooling		[183]
Adaraka	<i>Zingiber officinale</i> Rosc.	Zingiberaceae	Rhizome	Zingerone	Loperamide (5 mg/kg, i.p.)	Intraluminal pressure changes and expelled fluid volume from the colon	Inhibited spontaneous contractile movements in the isolated colonic segments, Inhibit colonic motility via direct action on smooth muscles	[184]
Badara	<i>Ziziphus jujuba</i> Mill.	Rhamnaceae	Leaf	Aqueous extract		Castor oil and magnesium sulfate-induced diarrhea models		[185]
Badara	<i>Ziziphus mauritiana</i>	Rhamnaceae	Root	Methanollic extract (25 and 50 mg/kg)	Diphenoxylate (2.5, 5 mg/kg) orally	Castor oil-induced diarrhea model and castor oil-induced fluid accumulation, spontaneous movement of the isolated rabbit jejunum, gastrointestinal transit time	An inhibition of the spontaneous penular movement of the isolated rabbit jejunum and inhibited acetylcholine-induced contraction of rat ileum	[186]

PGE₂: Prostaglandin E₂, CCB: Calcium channel blockade, EPEC: Enteropathogenic *Escherichia coli*, EIEC: Enteroinvasive *Escherichia coli*




Figure 1: Antidiarrheal medicinal plants




Figure 2: Distribution of plant parts investigated for antidiarrheal activity

Many phytoconstituents such as lupinofolin isolated from *Eriosema chinense*, -omoneukanrin B, dimethoxyflavone isolated from the stem bark of *Stereospermum kunthianum*, 6-(4-hydroxy-3-methoxyphenyl)-hexanoic acid, isovanillin, iso-acetovanillon from *Pycnocyclus spinosa* Decne. Ex Boiss., have been evaluated for anti-diarrheal activity. However, in the mentioned list of ayurvedic plants limited isolation of the active constituents have been done which accounts for the numerous scope in this area for analytical, pharmacognostical as well as pharmacological screening of the active principles from these plants. Some of the constituents such as kuryyam, koenimbine, koenine, piperine, and berberine are mentioned in the list with reported antidiarrheal activity [189-192].

Newer technologies such as in-silico, docking studies, interaction with enterotoxin from causative organism and nanotechnology

were also employed in the antidiarrheal agent research works [193,194]. However, unfortunately, such advanced techniques were not used for the above listed ayurvedic plants. However, a few clinical trials reveal that the plants acts via a number of mechanisms, i.e., anti-inflammatory, antisecretory antimicrobial effect against *V. cholerae* and enterotoxigenic *E. coli*, rotavirus, detoxification of toxins and constipate, adsorbent, providing a rich source of calories; antitomotility and antispasmodics effects [195].

CONCLUSION

The ethnomedicinal approach for diarrhea is a practical, cost-effective, and a logical for its treatment. Present data show that only a few isolated compounds from plants were investigated for antidiarrheal potential. Therefore, a significant research of chemical and biological properties of such less explored plants is still needed to determine their antidiarrheal efficacy which will possibly define their exact mechanism of actions.

REFERENCES

- Ghai OP, Paul VK, Arvind B. Ghai Essential Pediatrics. 7th ed. New Delhi: CBS Publishers & Distributors Pvt., Ltd.; 2009. p. 559-61.
- Schiller LR. Definitions, pathophysiology, and evaluation of chronic diarrhoea. Best Pract Res Clin Gastroenterol 2012;26:551-62.
- Whyte LA, Jenkins HR. Pathophysiology of diarrhoea. Paediatr Child Health 2012;22:443-7.
- Teke GN, Kuate JR, Ngouateu OB, Gatsing D. Antidiarrhoeal and antimicrobial activities of *Emilia coccinea* (Sims) G. Don extracts. J Ethnopharmacol 2007;112:278-83.
- Kitaoka M, Miyata ST, Unterweger D, Pukatzki S. Antibiotic resistance mechanisms of *Vibrio cholerae*. J Med Microbiol 2011;60:397-407.
- Baldi F, Bianco MA, Nardone G, Pilotto A, Zamparo E. Focus on acute diarrhoeal disease. World J Gastroenterol 2009;15:3341-8.
- Pimentel M, Hwang L, Melmed GY, Low K, Vasiliauskas E, Ippoliti A, *et al.* New clinical method for distinguishing D-IBS from other gastrointestinal conditions causing diarrhea: The LA/IBS diagnostic

- strategy. *Dig Dis Sci* 2010;55:145-9.
8. Al Jarousha AM, El Jarou MA, El Qouqa IA. Bacterial enteropathogens and risk factors associated with childhood diarrhea. *Indian J Pediatr* 2011;78:165-70.
 9. Tumwine JK, Thompson J, Katua-Katua M, Mujwajuzi M, Johnstone N, Porras I. Diarrhoea and effects of different water sources, sanitation and hygiene behaviour in East Africa. *Trop Med Int Health* 2002;7:750-6.
 10. Barkun AN, Love J, Gould M, Pluta H, Steinhart H. Bile acid malabsorption in chronic diarrhea: Pathophysiology and treatment. *Can J Gastroenterol* 2013;27:653-9.
 11. Page AL, Hustache S, Luquero FJ, Djibo A, Manzo ML, Grais RF. Health care seeking behavior for diarrhea in children under 5 in rural Niger: Results of a cross-sectional survey. *BMC Public Health* 2011;11:389.
 12. Rahman MK, Barua S, Islam MF, Islam MR, Sayeed MA, Parvin MS, *et al.* Studies on the anti-diarrheal properties of leaf extract of *Desmodium puchellum*. *Asian Pac J Trop Biomed* 2013;3:639-43.
 13. Casburn-Jones AC, Farthing MJ. Management of infectious diarrhoea. *Gut* 2004;53:296-305.
 14. Lamberti LM, Walker CL, Chan KY, Jian WY, Black RE. Oral zinc supplementation for the treatment of acute diarrhea in children: A systematic review and meta-analysis. *Nutrients* 2013;5:4715-40.
 15. Hickson M. Probiotics in the prevention of antibiotic-associated diarrhoea and *Clostridium difficile* infection. *Therap Adv Gastroenterol* 2011;4:185-97.
 16. Leung WK, Wu JC, Liang SM, Chan LS, Chan FK, Xie H, *et al.* Treatment of diarrhea-predominant irritable bowel syndrome with traditional Chinese herbal medicine: A randomized placebo-controlled trial. *Am J Gastroenterol* 2006;101:1574-80.
 17. Yamakawa J, Moriya J, Takeuchi K, Nakatou M, Motoo Y, Kobayashi J. Significance of Kampo, Japanese traditional medicine, in the treatment of obesity: Basic and clinical evidence. *Evid Based Complement Alternat Med* 2013;2013:943075.
 18. Manheimer E, Wieland LS, Cheng K, Li SM, Shen X, Berman BM, *et al.* Acupuncture for irritable bowel syndrome: Systematic review and meta-analysis. *Am J Gastroenterol* 2012;107:835-47.
 19. Mishra A, Sharma V, Hem K, Maurya SK. Plants used for treatment of diarrhea: An ayurvedic prospective. *Innov J Ayurvedic Sci* 2015;3:1-6.
 20. Kunal G, Sanjiv K, Vijayanand W. Anti diarrhoeal activity of a polyherbal formulation in various animals models diarrhoea. *Int Res J Pharm* 2012;3:289-90.
 21. Laloo D, Hemalatha S. Ethnomedicinal plants used for diarrhea by tribals of Meghalaya, Northeast India. *Pharmacogn Rev* 2011;5:147-54.
 22. Jachak SM, Saklani A. Challenges and opportunities in drug discovery from plants. *Curr Sci* 2007;92:1251-1.
 23. Gutiérrez SP, Sánchez MA, González CP, García LA. Antidiarrhoeal activity of different plants used in traditional medicine. *Afr J Biotechnol* 2007;6:2988-94.
 24. Komal, Kumar S, Rana AC. Herbal approaches for diarrhoea: A review. *Int Res J Pharm* 2013;4:31-8.
 25. Birdi TJ, Brijesh S, Daswani PG. Bactericidal effect of selected antidiarrhoeal medicinal plants on intracellular heat-stable enterotoxin-producing *Escherichia coli*. *Indian J Pharm Sci* 2014;76:229-35.
 26. Tetali P, Waghchaure C, Daswani PG, Antia NH, Birdi TJ. Ethnobotanical survey of antidiarrhoeal plants of Parinche valley, Pune district, Maharashtra, India. *J Ethnopharmacol* 2009;123:229-36.
 27. De Wet H, Nkwanyana MN, van Vuuren SF. Medicinal plants used for the treatment of diarrhoea in northern Maputaland, KwaZulu-Natal province, South Africa. *J Ethnopharmacol* 2010;130:284-9.
 28. Weiss RF. In: Arcanum AB, editor. *Herbal Medicine*. Beaconsfield, UK: Beaconsfield Publishers; 1998. p. 22-8.
 29. Chopra RN, Nayer SL, Chopra IC. *Glossary of Indian Medicinal Plants*. New Delhi: Council of Scientific and Industrial Research; 1956.
 30. Abubakar K, Abubakar MR, Ugwah-Oguejiofor JC, Muhammad AA, Usman M, Mshelia HE. Antidiarrhoeal activity of the saponin and flavonoid fractions of *Anarcadium occidentale* leaves in albino rats. *Adv Med Plant Res* 2015;3:23-8.
 31. Galvez J, Zarzuelo A, Crespo ME, Lorente MD, Ocete MA, Jiménez J. Antidiarrhoeal activity of *Euphorbia hirta* extract and isolation of an active flavonoid constituent. *Planta Med* 1993;59:333-6.
 32. Yadav AK, Tangpu V. Antidiarrheal activity of *Lithocarpus dealbata* and *Urena lobata* extracts. *Ther Implications Pharm Biol* 2007;45:223-9.
 33. Pérez GS, Pérez GC, Zavala MA. A study of the antidiarrheal properties of *Loeselia mexicana* on mice and rats. *Phytomedicine* 2005;12:670-4.
 34. Kumar R, Sharma RJ, Bairwa K, Roy RK, Kumar A. Pharmacological review on natural antidiarrhoeal agents. *Der Pharm Chem* 2010;2:66-93.
 35. Wang GS, Han J, Zhao LW, Jiang DX, Liu YT, Liu XL. Anthelmintic activity of steroidal saponins from *Paris polyphylla*. *Phytomedicine* 2010;17:1102-5.
 36. Dubreuil JD. Antibacterial and antidiarrheal activities of plant products against enterotoxigenic *Escherichia coli*. *Toxins* 2013;5:2009-41.
 37. Chandrashekar VM, Nagappa AN, Channesh TS, Habbu PV, Rao KP. Antidiarrhoeal activity of *Abutilon indicum* Linn leaf extracts. *J Nat Rem* 2004;4:12-6.
 38. Ray D, Sharatchandra KH, Thokchom IS. Antipyretic, antidiarrheal, hypoglycaemic and hepatoprotective activities of ethyl acetate extract of *Acacia catechu* Willd. in albino mice. *Ind J Pharmacol* 2006;38:408-13.
 39. Misra A, Bhagat R, Mujumdar AM. Antidiarrhoeal activity of *Acacia nilotica* Willd. bark methanol extract. *Hindustan Antibiot Bull* 2007-2008;49-50:14-20.
 40. Prasad SK, Jain D, Patel DK, Sahu AN, Hemalatha S. Antisecretory and antimotility activity of *Aconitum heterophyllum* and its significance in treatment of diarrhea. *Indian J Pharmacol* 2014;46:82-7.
 41. Rajput SB, Tonge MB, Karuppaiyil SM. An overview on traditional uses and pharmacological profile of *Acorus calamus* Linn. (Sweet flag) and other *Acorus* species. *Phytomedicine* 2014;21:268-76.
 42. Shoba FG, Thomas M. Study of antidiarrhoeal activity of four medicinal plants in castor-oil induced diarrhoea. *J Ethnopharmacol* 2001;76:73-6.
 43. Brijesh S, Daswani P, Tetali P, Antia N, Birdi T. Studies on the antidiarrhoeal activity of *Aegle marmelos* unripe fruit: Validating its traditional usage. *BMC Complement Altern Med* 2009;9:47.
 44. Rao GHJ, Lakshmi P. Evaluation of Antidiarrhoeal activity of extract from leaves of *Aegle marmelos*. *J App Pharm Sci* 2012;02:75-8.
 45. Shamkuwar PB, Sahi SR, Pawar DP. Efficacy of *Aegle marmelos* L. *Corr.* (Rutaceae) in Ayurvedic antidiarrhoeal formulation. *European J Exp Biol* 2012;2:194-8.
 46. Surve VS, Ghangale GR, Tamhankar SP, Gatne MM. Studies on antidiarrhoeal activity of *Aegle marmelos* (bael) in rats. *The J Bombay Vet Col* 2008;16:27-33.
 47. Rao CHV, Amresh KR, Irfan A, Rawat AKS, Pushpangadan P. Protective effect of *Aegle marmelos* fruit in gastrointestinal dysfunction in rats. *J Pharm Bio* 2003;41:558-63.
 48. Joshi PV, Patil RH, Maheshwari VL. *In vitro* antidiarrhoeal activity and toxicity profile of *Aegle marmelos* dried fruit pulp. *Nat Prod Radiance* 2009;8:498-502.
 49. Mazumder R, Bhattacharya S, Mazumder A, Pattnaik AK, Tiwary PM, Chaudhary S. Antidiarrhoeal evaluation of *Aegle Marmelos* (Correa) Linn. Root extract. *Phytother Res* 2006;20:82-4.
 50. Dhuley JN. Investigation on the gastroprotective and antidiarrhoeal properties of *Aegle marmelos* unripe fruit extract. *Hindustan Antibiot Bull* 2003-2004;45-46:41-6.
 51. Singh S, Bhandari A, Rai AK, Sharma P, Birshiliya Y, Gupta MK, *et al.* Comparative study on antidiarrhoeal effect and phytochemical screening of aqueous and alcoholic extract of *Aerva lanata*. *Asian J Chem* 2012;24:4545-47.
 52. Kumar N, Sharma SK. Evaluation of antidiarrhoeal potential of *Ailanthus excelsa* (Roxb) bark extract in rats. *Am Res J Pharm* 2015;1:1-4.
 53. Besra SE, Gomes A, Chaudhury L, Vedasiromoni JR, Ganguly DK. Antidiarrhoeal activity of seed extract of *Albizzia lebbbeck* Benth. *Phytother Res* 2002;16:529-33.
 54. Shah AJ, Gowani SA, Zuberi AJ, Ghayur MN, Gilani AH. Antidiarrhoeal and spasmolytic activities of the methanolic crude extract of *Alstonia scholaris* L. are mediated through calcium channel blockade. *Phytother Res* 2010;24:28-32.
 55. Saifuzzaman, Hossain A, ShahinUA, Islam A, Ali ES. Antidiarrheal and cytotoxic activities of *Alstonia scholaris* bark. *Int Res J Pharm* 2013;4:101-3.
 56. Hussain Z, Amresh G, Singh S, Rao CV. Antidiarrheal and antiulcer activity of *Amaranthus spinosus* in experimental animals. *Pharm Bio* 2009;47:932-9.
 57. Dharmalingam SR, Madhappan R, Ramamurthy S, Chidambaram K, Srikanth MV, Shanmugham S, *et al.* Investigation on antidiarrhoeal activity of *Aristolochia indica* Linn. Root extracts in mice. *Afr J Tradit Complement Altern Med* 2014;11:292-4.

58. Venkatesan N, Thiyagarajan V, Narayanan S, Arul A, Raja S, Vijaya Kumar SG, et al. Anti-diarrhoeal potential of *Asparagus racemosus* wild root extracts in laboratory animals. *J Pharm Pharm Sci* 2005;8:39-46.
59. Sandhyarani G, Swathi K, Gangarapu K, Kumar PK, Ramesh A. Pharmacological studies of anti-diarrhoeal activity of *Barringtonia acutangula* (L.) In experimental animals. *Int J Phyto Pharm Res* 2014;5:93-5.
60. Zafar Imam M, Sultana S, Akter S. Antinociceptive, antidiarrheal, and neuropharmacological activities of *Barringtonia acutangula*. *Pharm Biol* 2012;50:1078-84.
61. Mukherjee KPK, Gopal TK, Subburaju T, Dhanbal SB, Duraiswamy B, Elango K, et al. Studies on the antidiarrheal profiles of *Bauhinia purpurea* L. Leaves (*Caesalpiniaaceae*) extract. *Nat Prod Sci* 1998;4:234-7.
62. Sengupta R, Ahmed AB. Evaluation of antidiarrheal activity of ethanolic extract of *Bauhinia variegata* (Leguminosae) stem bark in Wistar albino rats. *Int J Pharm Biomed Res* 2015;2015:201503.
63. Shamkuwar PB, Pawar DP. Antidiarrhoeal and antispasmodic effect of *Berberis aristata*. *Int J Pharmacogn Phytochem Res* 2013;5:24-6.
64. Joshi PV, Shirkhedkar AA, Prakash K, Maheshwari VL. Antidiarrheal activity, chemical and toxicity profile of *Berberis aristata*. *Pharm Biol* 2011;49:94-100.
65. Sheikh NW, Upwar NI, Patel RD, Rupapara DJ. Evaluation of antidiarrhoeal activity of alcoholic extract of stem of *Berberis aristata* DC. *J Pharm Res* 2010;3:2232-4.
66. Shilpi JA, Taufiq-ur-Rahman M, Uddin SJ, Alam MS, Sadhu SK, Seidel V. Preliminary pharmacological screening of *Bixa orellana* L. Leaves. *J Ethnopharmacol* 2006;108:264-71.
67. Borrelli F, Capasso F, Capasso R, Ascione V, Aviello G, Longo R, et al. Effect of *Boswellia serrata* on intestinal motility in rodents: Inhibition of diarrhoea without constipation. *Br J Pharmacol* 2006;148:553-60.
68. Sharma U, Lahkar M, Lahon J. Evaluation of antidiarrhoeal potential of *Bryophyllum Pinnatum* in experimental animals. *Asian J Biomed Pharm Sci* 2012;2:28-31.
69. Sumithra M, Anbu J, Suganya S. Antidiarrheal activity of the methanolic extract of *Bunchaniana lanzan* against castor oil induction. *Int J Adv Pharm Bio Chem* 2012;1:151-3.
70. Gunakkunru A, Padmanaban K, Thirumal P, Pritila J, Parimala G, Vengatesan N, et al. Antidiarrhoeal activity of *Butea monosperma* in experimental animals. *J Ethnopharmacol* 2005;98:241-4.
71. Billah MM, Islam R, Khatun H, Parvin S, Islam E, Islam SA, et al. Antibacterial, antidiarrhoeal, and cytotoxic activities of methanol extract and its fractions of *Caesalpinia bonducella* (L.) Roxb leaves. *BMC Complement Altern Med* 2013;13:101.
72. Afroz T, Ramproshad S, Mondal B, Haque A, Khan R. Antidiarrhoeal and analgesic activity of barks of medicinal plant *Caesalpinia pulcherrima*. *Int J Pharm Sci Res* 2013;4:1946-9.
73. Havagiray R, Ramesh C, Sadhna K. Study of antidiarrhoeal activity of *Calotropis gigantea* R. Br. in experimental animals. *J Pharm Sci* 2004;7:70-5.
74. Kumar S, Dewan S, Sangraula H, Kumar VL. Anti-diarrhoeal activity of the latex of *Calotropis procera*. *J Ethnopharmacol* 2001;76:115-8.
75. Das A, Dutta AK, Razzaque S, Saha B, Gope PS, Choudhury N. Analgesic and antidiarrheal properties of the latex of *Calotropis Procera*. *Int J Pharm Bio Arch* 2011;2:521-5.
76. Patil SH, Adkar PP, Shelke TT, Oswal RJ, Borase SP. Antidiarrhoeal activity of methanol extract of leaves of *Calotropis procera* R. Br. *J Pharm Toxicol* 2011;1:25-30.
77. Abhinayani G, Sravya N, Naga Kishore R. Anti-diarrheal activity of alcoholic and aqueous extract of *Calotropis procera* R. Br. leaves in rats. *Int J Pharm Pharm Sci* 2013;5:878-80.
78. Besra SE, Gomes A, Ganguly DK, Vedasiromoni JR. Antidiarrhoeal activity of hot water extract of black tea (*Camellia sinensis*). *Phytother Res* 2003;17:380-4.
79. Sini KR, Sinha BN, Rajasekaran A. Antidiarrheal activity of *Capparis zeylanica* leaf extracts. *J Adv Pharm Technol Res* 2011;2:39-42.
80. Mahboob AS, Gouda TS. A study on anti-diarrhoeal activity of fruit extracts of *Carica papaya* (Caricaceae) Linn. in rat. *Pharmatutor art* 1372. Available from: <http://www.pharmatutor.org/articles/study-anti-diarrhoeal-activity-fruit-extracts-carica-papaya-caricaceae-linn-rats>. [Last accessed on 2015 Sep 26].
81. Sharma P, Vidyasagar G, Singh S, Ghule S, Kumar B. Antidiarrhoeal activity of leaf extract of celosia argentea in experimentally induced diarrhoea in rats. *J Adv Pharm Technol Res* 2010;1:41-8.
82. Rao CV, Vijayakumar M, Sairam K, Kumar V. Antidiarrhoeal activity of the standardised extract of *Cinnamomum tamala* in experimental rats. *J Nat Med* 2008;62:396-402.
83. Rao HJ, Lakshmi. Antidiarrhoeal activity of the aqueous extract of the bark of *Cinnamomum zeylanicum* Linn in mice. *J Clin Diagn Res* 2012;6:215-9.
84. Amresh, Reddy GD, Rao CV, Shirwaikar A. Ethnomedical value of *Cissampelos pareira* extract in experimentally induced diarrhoea. *Acta Pharm* 2004;54:27-35.
85. Devi BP, Boominathan R, Mandal SC. Evaluation of anti-diarrheal activity of *Cleome viscosa* L. extract in rats. *Phytomedicine* 2002;9:739-42.
86. Pal A, Mahmud ZA, Akter N. Evaluation of antinociceptive, antidiarrheal and antimicrobial activities of leaf extracts of *Clerodendrum indicum*. *Pharmacogn J* 2012;4:41-6.
87. Sini KR, Rajasekaran A, Sangeetha PT. Anti-diarrheal activity of the leaves of *Clitoria ternatea* L. *Int Res J Pharm Sci* 2011;2:7-9.
88. Upwar NK, Patel R, Waseem N, Mahobia NK. Evaluation of antidiarrhoeal activity of the root of *Clitoria ternatea* Linn. *Int J Pharm Sci Rev Res* 2010;5:131-4.
89. Nithya V. Evaluation of antidiarrheal activity on *Coriandrum sativum* Linn. in wistar albino rats. *World J Pharm Res* 2015;4:638-43.
90. Inayathulla, Shariff WR, Karigar AA, Sikarwar MS. Evaluation of anti diarrhoeal activity of *Crataeva nurvala* root bark in experimental animals. *Int. J Pharm Pharm Sci* 2010;2:158-61.
91. Sahoo HB, Sahoo SK, Sarangi SP, Sagar R, Kori ML. Anti-diarrhoeal investigation from aqueous extract of *Cuminum cyminum* Linn. Seed in Albino rats. *Pharmacognosy Res* 2014;6:204-9.
92. Owolabi OJ, Arhewoh MI, Aadum EJ. Evaluation of the antidiarrhoeal activity of the aqueous rhizome extract of *Curcuma Longa*. *J Pharm Allied Sci* 2012;9:1450-7.
93. Ravindra BDS, Neeharika V, Pallavi V, Reddy MB. Antidiarrheal activity of *Cynodon Dactylon*. *Pers. Phcog Mag* 2009;5:23-7.
94. Uddin SJ, Mondal K, Shilpi JA, Rahman MT. Antidiarrhoeal activity of *Cyperus rotundus*. *Fitoterapia* 2006;77:134-6.
95. Daswani PG, Brijesh S, Tetali P, Birdi TJ. Studies on the activity of *Cyperus rotundus* Linn. Tubers against infectious diarrhea. *Indian J Pharmacol* 2011;43:340-4.
96. Mujumdar AM, Misar AV, Upadhye AS. Antidiarrhoeal activity of ethanol extract of the bark of *Dalbergia lanceolaria*. *J Ethnopharmacol* 2005;102:213-6.
97. Chandra P, Sachan N, Pal D. Protective effect of *Dalbergia sissoo* Roxb. Ex DC. (family: Fabaceae) leaves against experimentally induced diarrhoea and peristalsis in mice. *Toxicol Ind Health* 2015;31:1229-35.
98. Medha MH, Lakshman K, Girija K, Kumar BA, Lakshmi prasanna V. Assessment of antidiarrhoeal activity of *Desmostachya bipinnata* L. (*Poaceae*) root extracts. *Bol Latinoam Caribe Plant Med Aromat* 2010;9:312-8.
99. Jayakumari S, Srinivas RGH, Anbu J, Ravichandiran V. Antidiarrheal activity of *Dichrostachys cinera* (L.) Weight & Arn. *Int J Pharm Sci* 2011;3:61-3.
100. Rouf R, Uddin SJ, Shilpi JA, Toufiq-ur-Rahman M, Ferdous MM, Sarker SD. Anti-diarrhoeal properties of *Diospyros peregrina* in the castor oil-induced diarrhoea model in mice. *Ars Pharm* 2006;47:81-9.
101. Hossen SMM, Sarkar R, Mahmud S, Aziz NMA. Medicinal potential of *Phyllanthus emblica* (Linn.) Fruits extracts: Biological and pharmacological activities. *Br J Pharm Res* 2014;4:1486-99.
102. Mehmood MH, Siddiqi HS, Gilani AH. The antidiarrheal and spasmolytic activities of *Phyllanthus emblica* are mediated through dual blockade of muscarinic receptors and Ca²⁺ channels. *J Ethnopharmacol* 2011;133:856-65.
103. Perianayagam JB, Narayanan S, Gnanasekar G, Pandurangan A, Raja S, Rajagopal K, et al. Evaluation of antidiarrheal potential of *Embllica officinalis*. *Pharm Bio* 2005;43:373-7.
104. Kamalraj R. Anti-diarrhoeal potential of *Erythrina indica* Lam – Leaf extracts in laboratory animals. *Int J Pharm Sci Drug Res* 2011;3:155-7.
105. Khan IN, Jahan S, Bhuiya MAM, Mazumder K, Saha BK. Anti-diarrheal potential of ethanol and water extracts of *Euphorbia hirta* whole plant on experimental animals: A comparative study. *Sch J Appl Med Sci* 2013;1:199-204.
106. Bellah SF, Raju MI, Billah SM, Rahman SE, Murshid GM, Rahman MM. Evaluation of antibacterial and antidiarrhoeal activity of ethanolic extract of *Feronia limonia* leaves. *Pharm Innov J* 2015;3:50-4.

107. Momin MAM, Khan MR, Rayhan J, Afrose A, Rana S, Begum AA. Evaluation of antibacterial and antidiarrhoeal activities of *Feronia limonia* leaf extract. *Am J Plant Sci* 2013;4:2181-85.
108. Mahalakshmi M, Parimala M, Shoba FG. Evaluation of anti-diarrhoeal potential of methanol extract of *Ficus bengalensis* Linn. leaf and *Mangifera indica* Linn. Stem bark and root bark. *Int J Pharmacogn Phytochem Res* 2014;6:454-8.
109. Mukherjee PK, Saha K, Murugesan T, Mandal SC, Pal M, Saha BP. Screening of anti-diarrhoeal profile of some plant extracts of a specific region of West Bengal, India. *J Ethnopharmacol* 1998;60:85-9.
110. Mandal SC, Ashok Kumar CK. Studies on anti-diarrhoeal activity of *Ficus hispida*. Leaf extract in rats. *Fitoterapia* 2002;73:663-7.
111. Panchawat S, Sisodia SS. Evaluation of anti-diarrhoeal activity of stem bark extracts of *Ficus religiosa* prepared by different methods of extraction. *Int J Pharm Biol Arch* 2012;3:218-22.
112. Pampattiwar SP, Advani NV. Evaluation of anti-diarrhoeal activity of *Ficus glomerata* in castor oil induced diarrhoea in rats. *J Sci* 2011;1:26-30.
113. Rehman N, Bashir S, Al-Rehaily AJ, Gilani AH. Mechanisms underlying the antidiarrheal, antispasmodic and bronchodilator activities of *Fumaria parviflora* and involvement of tissue and species specificity. *J Ethnopharmacol* 2012;144:128-37.
114. Panda SK, Das D, Tripathy NK. Antidiarrhoeal activity of various root extracts of *Gmelina arborea* roxb. In experimentally induced diarrhoea in mice. *World J Pharm Pharm Sci* 2015;4:912-9.
115. Agunu A, Yusuf S, Andrew GO, Zezi AU, Abdurahman EM. Evaluation of five medicinal plants used in diarrhoea treatment in Nigeria. *J Ethnopharmacol* 2005;101:27-30.
116. Das S, Prakash R, Devaraj SN. Antidiarrhoeal effects of methanolic root extract of *Hemidesmus indicus* (Indian sarsaparilla) – An *in vitro* and *in vivo* study. *Indian J Exp Biol* 2003;41:363-6.
117. Shalini R, Rajan S. Antidiarrhoeal activity of aqueous and alcoholic extracts of *Hemidesmus indicus* Root. *Int J Pharm Pharm Sci* 2015;7:403-6.
118. Kavitha D, Shilpa PN, Devaraj SN. Antibacterial and antidiarrhoeal effects of alkaloids of *Holarrhena antidysenterica* wall. *Indian J Exp Biol* 2004;42:589-94.
119. Sharma S, Lakshmi KS, Rajesh T. Evaluation of antidiarrhoeal potentials of ethanolic extract of leaves of *Holoptelea integrifolia* in mice model. *Int J Pharm Tech Res* 2009;1:832-6.
120. Maniyar Y, Bhixavatimath P, Agashikar NV. Antidiarrhoeal activity of flowers of *Ixora coccinea* Linn. in rats. *J Ayurved Integr Med* 2010;1:287-91.
121. Yasmeen M, Prabhu B, Agashikar NV. Evaluation of the anti-diarrhoeal activity of the leaves of *Ixora coccinea* Linn. in rats. *J Clin Diagn Res* 2010;4:3298-303.
122. Mujumdar AM, Misar AV, Salaskar MV, Upadhye AS. Antidiarrhoeal effect of an isolated fraction (JC) of *Jatropha curcas* roots in mice. *J Nat Rem* 2001;1/2:89-93.
123. Mujumdar AM, Upadhye AS, Misar AV. Studies on antidiarrhoeal activity of *Jatropha curcas* root extract in albino mice. *J Ethnopharmacol* 2000;70:183-7.
124. Rahman MD, Haque ME, Solaiman M, Saifuzzaman M. Atiociceptive and antidiarrhoeal activities of *Madhuca idica* J.F.Gmel. *Pharmacologyonline* 2011;1:473-80.
125. Rajan S, Suganya H, Thirunalasundari T, Jeeva S. Antidiarrhoeal efficacy of *Mangifera indica* seed kernel on Swiss albino mice. *Asian Pac J Trop Med* 2012;5:630-3.
126. Yakubu MT, Salimon SS. Antidiarrhoeal activity of aqueous extract of *Mangifera indica* L. Leaves in female albino rats. *J Ethnopharmacol* 2015;163:135-41.
127. Sairam K, Hemalatha S, Kumar A, Srinivasan T, Ganesh J, Shankar M, *et al.* Evaluation of antidiarrhoeal activity in seed extracts of *Mangifera indica*. *J Ethnopharmacol* 2003;84:11-5.
128. Shah AJ, Bhulani NN, Khan SH, Ur Rehman N, Gilani AH. Calcium channel blocking activity of *Mentha longifolia* L. Explains its medicinal use in diarrhoea and gut spasm. *Phytother Res* 2010;24:1392-7.
129. Jalilzadeh-Amin G, Maham M. Antidiarrheal activity and acute oral toxicity of *Mentha longifolia* L. Essential oil. *Avicenna J Phytomed* 2015;5:128-37.
130. Khalid MS, Kumar SJ, Suresh DK, Singh RJ, Reddy IVN, Kumar S. Evaluation of anti-diarrhoeal potential of ethanolic extract of *Mimosa pudica* leaves. *Int J Green Pharm* 2011;5:75-8.
131. Balakrishnan N, Suresh D, Pandian GS, Edwin E, Sheeja E. Antidiarrhoeal potential of *Mimosa pudica* root extracts. *Indian J Nat Prod* 2006;22:21-3.
132. Bakare RI, Magbagbeola OA, Akinwande AI, Okunowo OW, Green M. Antidiarrhoeal activity of aqueous leaf extract of *Momordica charantia* in rats. *J Pharmacognosy Phytother* 2011;3:1-7.
133. Lakshminarayana M, Shivkumar H, Rimaben P, Bhargava VK. Antidiarrhoeal activity of leaf extract of *Moringa oleifera* in experimentally induced diarrhoea in rats. *Int J Phytomed* 2011;3:68-74.
134. Misra A, Srivastava S, Srivastava M. Evaluation of anti diarrheal potential of *Moringa oleifera* (Lam.) leaves. *J Pharmacogn Phytochem* 2014;2:43-6.
135. Sharma P, Vidyasagar G, Bhandari A, Singh S, Bhadoriya U, Ghule S, *et al.* Pharmacological evaluation of antidiarrhoeal activity of leaves extract of *Murraya koenigii* in experimentally induced diarrhoea in rats. *Asian Pac J Trop Dis* 2012;2:230-3.
136. Mandal S, Nayak A, Kar M, Banerjee SK, Das A, Upadhyay SN, *et al.* Antidiarrhoeal activity of carbazole alkaloids from *Murraya koenigii* Spreng (*Rutaceae*) seeds. *Fitoterapia* 2010;81:72-4.
137. Rahman MD, Hasanuzzaman M, Uddin N, Shahid IZ. Anti-diarrhoeal and anti-inflammatory activities of *Murraya pauculata* (L.) Jack. *Pharmacologyonline* 2010;3:768-76.
138. Yakubu MT, Nurudeen QO, Salimon SS, Yakubu MO, Jimoh RO, Nafiu MO, *et al.* Antidiarrhoeal activity of *Musa paradisiaca* Sap in Wistar rats. *Evid Based Complement Alternat Med* 2015;2015:683726.
139. Grover JK, Khandkar S, Vats V, Dhunoo Y, Das D. Pharmacological studies on *Myristica fragrans* – Antidiarrheal, hypnotic, analgesic and hemodynamic (blood pressure) parameters. *Methods Find Exp Clin Pharmacol* 2002;24:675-80.
140. Talukder MJ, Nessa J. Effect of *Nelumbo nucifera* rhizome extract on the gastrointestinal tract of rat. *Bangladesh Med Res Council Bull* 1998;24:6-9.
141. Mukherjee PK, Das J, Balasubramanian R, Saha K, Pal M, Saha BP. Antidiarrhoeal evaluation of *Nelumbo nucifera* rhizome extract. *Indian J Pharmacol* 1995;27:262-4.
142. Arumugam A, Dhailappan AK. Fatty acid composition and antidermatophytic and antidiarrheal activity of *Nelumbo nucifera* seed oil. *Int J Pharm Pharm Sci* 2012;4:769-75.
143. Shareef H, Rizwani GH, Mandukhail SR, Watanabe N, Gilani AH. Studies on antidiarrhoeal, antispasmodic and bronchodilator activities of *Operculina turpethum* Linn. *BMC Complement Altern Med* 2014;14:479.
144. Asaduzzaman M, Nasrin N, Muhit MA, Raihan SZ, Apu AS, Akbar. Antidiarrheal, analgesic and Cytotoxic activities of crude extract of *Oroxylum indicum* (L.) stem bark. *J Pharm Res* 2011;4:4296-98.
145. Joshi SV, Gandhi TR, Vyas BA, Shah PD, Patel PK, Vyas HG. Effect of *Oroxylum indicum* on intestinal motility in rodents. *Orient Pharm Exp Med* 2012;12:279-85.
146. Pierre W, Evelyne N, Telesphore B, Nguetefack TB, Wansi SL, Albert K. Antidiarrhoeal efficacy of aqueous and methanolic extracts of *Oxalis corniculata* Klotzsch. in rats. *Cameroon J Exp Bio* 2005;1:46-9.
147. Afroz S, Alamgir M, Khan MT, Jabbar S, Nahar N, Choudhuri MS. Antidiarrhoeal activity of the ethanol extract of *Paederia foetida* Linn. (*Rubiaceae*). *J Ethnopharmacol* 2006;105:125-30.
148. Abel NA, Helen OK, Wilson OH. Antidiarrheal activity of aqueous fruit extract of *Phoenix dactylifera* (date palm) in wistar rats. *Bri J Pharmacol Toxicol* 2013;4:121-7.
149. Shamkuwar PB, Shahi SR. Study of antidiarrhoeal activity of piperine. *Der Pharmacia Lettre* 2012;4:217-21.
150. Shamkuwar PB, Shahi SR, Jadhav ST. Evaluation of antidiarrhoeal effect of black pepper (*Piper nigrum* L.). *Asian J Plant Sci Res* 2012;2:48-53.
151. Shamkuwar PB. Mechanisms of antidiarrhoeal effect of *Piper nigrum*. *Int J Pharm Tech Res* 2013;5:1138-41.
152. Bajad S, Bedi KL, Singla AK, Johri RK. Antidiarrhoeal activity of piperine in mice. *Planta Med* 2001;67:284-7.
153. Sih T, Shah AJ, Gilani AH. Insight into the possible mechanism of antidiarrheal and antispasmodic activities of piperine. *Pharm Biol* 2009;47:660-4.
154. Janbaz KH, Hassan W, Mehmood MH, Gilani AH. Antidiarrheal, antispasmodic and bronchodilator activities of *Pistacia integerrima* are mediated through dual inhibition of muscarinic receptors and Ca⁺⁺ influx. *Sci Tech Dev* 2015;34:52-9.
155. Brijesh S, Daswani PG, Tetali P, Rojatkhar SR, Antia NH, Birdi TJ. Studies on *Pongamia pinnata* L. Pierre leaves: Understanding the

- mechanism(s) of action in infectious diarrhea. J Zhejiang Univ Sci B 2006;7:665-74.
156. Ojewole JA, Awe EO, Chiwororo WD. Antidiarrhoeal activity of *Psidium guajava* Linn. (*Myrtaceae*) leaf aqueous extract in rodents. J Smooth Muscle Res 2008;44:195-207.
 157. Birdi T, Daswani P, Brijesh S, Tetali P, Natu A, Antia N. Newer insights into the mechanism of action of *Psidium guajava* L. leaves in infectious diarrhoea. BMC Complement Altern Med 2010;10:33.
 158. Prakash T, Bhoumik D, Shivakumar H, Venkatesh JS, Dutta AS. Antidiarrhoeal activity of *Psidium guajava* bark extracts. J Nat Rem 2007;7:278-82.
 159. Jai D, Patel I, Somai R. Anti-diarrhoeal activity of ethaolic heartwood extract of *Pterocarpus marsupium*. Pharmacologyonline 2011;1:552-9.
 160. Das AK, Mandal SC, Banerjee SK, Sinha S, Das J, Saha BP, et al. Studies on antidiarrhoeal activity of *Punica granatum* seed extract in rats. J Ethnopharmacol 1999;68:205-8.
 161. Qnais EY, Elokda AS, Ghalyun YYA, Abdulla FA. Antidiarrheal activity of the aqueous extract of *Punica granatum* (Pomegranate) peels. Pharm Biol 2007;45:715-20.
 162. Shamkuwar PB, Pawar DP. Evaluation of antidiarrhoeal potential of *Punica granatum* L. (Punicaceae) in Ayurvedic formulation. J Chem Pharm Res 2012;4:1489-92.
 163. Choudhary GP. Antidiarrhoeal activity of ethanolic extract of *Quercus infectoria*. Int J Pharm Chem Sci 2012;1:1404-7.
 164. Ezeigbo II, Ezeja MI, Madubuike KG, Ifenkwe DC, Ukwani IA, Udeh NE, et al. Antidiarrhoeal activity of leaf methanolic extract of *Rauwolfia serpentina*. Asian Pac J Trop Biomed 2012;2:430-2.
 165. Harde MT, Khairnar AS, Kastureand AS, Kasture SB. Evaluation of antipsychotic and anti-diarrhoeal activities of ethanolic extract of roots of *Rubia cordifolia* Linn. Orient Pharm Exp Med 2008;8:73-80.
 166. Guo H, Zhang J, Gao W, Qu Z, Liu C. Anti-diarrhoeal activity of methanol extract of *Santalum album* L. In mice and gastrointestinal effect on the contraction of isolated jejunum in rats. J Ethnopharmacol 2014;154:704-10.
 167. Panchawat S, Sisodia SS. *In-vivo* antidiarrhoeal activity of extracts from stem bark of *Saraca asoca* Roxb. Prepared by different extraction methods. Intl J Pharm Bio Sci 2012;2:338-43.
 168. Negi JS, Bisht VK, Bhandari AK, Bhatt VP, Sati MK, Mohanty JP, et al. Antidiarrheal activity of methanol extract and major essential oil contents of *Saussurea lappa* Clarke. Afr J Pharm Pharmacol 2013;7:474-7.
 169. Sarangi RR, Mishra US, Panda SK, Behera S. Evaluation of antidiarrhoeal activity of *Sida rhombifolia* Linn. Root. Int Res J Pharm 2011;2:157-60.
 170. Biswas S, Murugesan T, Sinha S, Maiti K, Gayen JR, Pal M, et al. Antidiarrhoeal activity of *Strychnos potatorum* seed extract in rats. Fitoterapia 2002;73:43-7.
 171. Mehjabeen, Ahmad M, Jahan N, Mahayrookh, Rehman AB, Muhammad S, et al. Antidiarrhoeal, anti-inflammatory and analgesic activities of *Symplocos racemosa* roxb. Bark. Pak J Pharm Sci 2014;27:2221-6.
 172. Shamkuwar PB, Pawar DP, Chauhan SS. Antidiarrhoeal activity of seeds of *Syzygium cumini* L. J Pharm Res 2012;5:5537.
 173. Janbaz KH, Qadir MI, Jan A, Gilani AH. Anti-diarrheal activity of methanolic extract of *Tephrosia purpurea*. Acta Pol Pharm 2013;70:345-7.
 174. Sivakumar G, Rao CG, Priya ESS, Shraavan N, Somasekhar P. Evaluation of anti-diarrhoeal activity of the bark of *Terminalia arjuna* [Roxb]. Int J Res Phytochem Pharmacol 2011;1:161-4.
 175. Bimelsh K, Kalyani D, Prashant T, Manoj S, Diwakar G. Evaluation of anti-diarrheal effect aqueous and ethanolic extracts of fruit pulp of *Terminalia bellerica* in rats. Int J Drug Dev Res 2010;2:769-79.
 176. Viswanatha GL, Hanumanthappa S, Krishnadas N, Rangappa S. Antidiarrheal effect of fractions from stem bark of *Thespesia populnea* in rodents: Possible antimotility and antisecretory mechanisms. Asian Pac J Trop Med 2011;4:451-6.
 177. Viswanatha GL, Srinath R, Nandakumar K, Shylaja H, Lakshman K. Antidiarrheal activity of alcoholic and aqueous extracts of stem bark of *Thespesia populnea* in rodents. Pharmacologyonline 2007;3:222-30.
 178. Kaur M, Singh A, Kumar B. Comparative antidiarrheal and antiulcer effect of the aqueous and ethanolic stem bark extracts of *Tinospora cordifolia* in rats. J Adv Pharm Technol Res 2014;5:122-8.
 179. Perianayagam JB, Sharma SK, Pillai KK. Evaluation of antidiarrheal potential of trichodesma indicum root extract in rats. Methods Find Exp Clin Pharmacol 2005;27:533-7.
 180. Boyina R, Kosanam S, Rani TT. Evaluation of anti-diarrheal activity of aqueous extract of *Trigonella foenum-graecum*. Int J Pharmacol Res 2014;4:130-3.
 181. Bashir S, Memon R, Gilani AH. Antispasmodic and antidiarrheal activities of valeriana hardwickii wall. Rhizome are putatively mediated through calcium Channel blockade. Evid Based Complement Altern Med 2011;2011:304960.
 182. Rajput MS, Nair V, Chauhan A, Jawanjal H, Dange VS. Evaluation of antidiarrheal activity of aerial parts of *Vinca major* in experimental animals. Middle East J Sci Res 2011;7:784-8.
 183. Bigoniya P, Rana AC. Antidiarrheal and antispasmodic activity of *Wrightia tinctoria* bark and its steroidal alkaloid fraction. Pharmacologyonline 2009;3:298-310.
 184. Iwami M, Shiina T, Hirayama H, Shima T, Takewaki T, Shimizu Y. Inhibitory effects of zingerone, a pungent component of *Zingiber officinale* Roscoe, on colonic motility in rats. J Nat Med 2011;65:89-94.
 185. Rao GH, Lakshmi P. Anti diarrhoeal activity of *Ziziphus jujuba* leaf extract in rats. Int J Pharm Bio Sci 2012;3:532-8.
 186. Dahiru D, Sini JM, John-Africa L. Antidiarrhoeal activity of *Ziziphus mauritiana* root extract in rodents. Afr J Biotechnol 2006;5:941-5.
 187. Jadavji TA. Caraka Samhita of Agnivesa, Elaborated by Caraka and Drinhbala with the Ayurveda Dipika Commentry by Cakrapanidatta. Varanasi: Chaukhamba Vidyabhawan; 2011.
 188. Sharma PV. Susruta Samhita, with English Translation of Text and Dalhana Commentary Along With Critical Notes. 1st ed. Varanasi: Chaukhamba Bharati Academy; 2001.
 189. Prasad SK, Laloo D, Kumar M, Hemalatha S. Antidiarrhoeal evaluation of root extract, its bioactive fraction, and lupinifolin isolated from *Eriosema chinense*. Planta Med 2013;79:1620-7.
 190. Ching FP, Otokitio IO, Egert-Omoneukanrin B. Dimethoxyflavone isolated from the stem bark of *Stereospermum kunthianum* possesses antidiarrhoeal activity in rodents. Afr J Tradit Complement Altern Med 2013;10:47-51.
 191. Sadraei H, Ghanadian M, Asghari G, Madadi E, Azali N. Antispasmodic and antidiarrhoeal activities of 6-(4-hydroxy-3-methoxyphenyl)-hexanoic acid from *Pycnocycla spinosa* Decne. Exboiss. Res Pharm Sci 2014;9:279-86.
 192. Sadraei H, Ghanadian M, Asghari G, Azali N. Antidiarrheal activities of isovanillin, iso-acetovanillon and *Pycnocycla spinosa* Decne ex.Boiss extract in mice. Res Pharm Sci 2014;9:83-9.
 193. Chen JC, Ho TY, Chang YS, Wu SL, Li CC, Hsiang CY. Identification of *Escherichia coli* enterotoxin inhibitors from traditional medicinal herbs by in silico, in vitro, and in vivo analyses. J Ethnopharmacol 2009;121:372-8.
 194. Velázquez C, Correa-Basurto J, Garcia-Hernandez N, Barbosa E, Tesoro-Cruz E, Calzada S, et al. Anti-diarrheal activity of (-)-epicatechin from *Chiranthodendron pentadactylon* Larreat: Experimental and computational studies. J Ethnopharmacol 2012;143:716-9.
 195. Asgari Z, Selwyn BJ, Vonville H, DuPont HL. A systematic review of the Evidence for use of herbal medicine for the treatment of acute diarrhea. Nat Prod J 2012;2:1-81.

© SAGEYA. This is an open access article licensed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted, noncommercial use, distribution and reproduction in any medium, provided the work is properly cited.

Source of Support: Nil, Conflict of Interest: None declared.