

REVIEW

doi: 10.5455/medarch.2018.72.214-219

MED ARCH. 2018 JUN; 72(3): 214-219

RECEIVED: MAR 10, 2018 | ACCEPTED: MAY 15, 2018

¹Department of Neurosurgery, Kermanshah University of Medical Sciences, Kermanshah, Iran²Medical Biology Research Center, Kermanshah University of Medical Sciences, Kermanshah, Iran³Students Research Committee, Kermanshah University of Medical Sciences, Kermanshah, Iran**Corresponding author:** Masoud Sadeghi, Medical Biology Research Center, Kermanshah University of Medical Sciences, Kermanshah, Iran. ORCID ID: <http://www.orcid.org/0000-0002-3586-3012>. E-mail: sadeghi_mbrc@yahoo.com

Development of Posttraumatic Hydrocephalus Requiring Ventriculoperitoneal Shunt After Decompressive Craniectomy for Traumatic Brain Injury: a Systematic Review and Meta-analysis of Retrospective Studies

Reza Fattahian¹, Seyed Reza Bagheri¹, Masoud Sadeghi^{2,3}**ABSTRACT**

Background: Decompressive craniotomy (DC) is a known risk factor for the development of posttraumatic hydrocephalus (PTH) in the patients with traumatic brain injury (TBI). Herein, the present study reported the development of PTH requiring ventriculoperitoneal (VP) shunt after DC for TBI. **Methods:** Four databases (PubMed, Web of Science, Scopus, and Cochrane Library) were searched from 1983 to April 2018. The studies evaluating the prevalence of PTH requiring VP shunt after DC in the patients with TBI were selected without language restriction. A random-effects meta-analysis using event rate (ER) and 95% confidence intervals (CIs), was run by RevMan5.3 software. **Results:** Out of 355 studies obtained from the databases, 25 studies were included and analyzed in the meta-analysis. The studies included 2402 patients undergoing DC for TBI, 354 of who had PTH. The pooled ER of hydrocephalus in the patients undergoing DC for TBI was 17.7% [95%CI: 13.2 to 23.4%; $P < 0.001$]. In addition, the pooled analysis showed that ER of hydrocephalus was 13% in adults [95%CI: 9 to 18.5%; $P < 0.001$] and 37.6% in children [95%CI: 27.79 to 48.7%; $P = 0.029$; $I^2 = 0\%$]. **Conclusion:** The present study demonstrated that DC after TBI was associated with the development of PTH, especially in children compared to adults.

Keywords: Traumatic brain injury, decompressive craniotomy, hydrocephalus, ventriculoperitoneal shunt.

1. INTRODUCTION

Traumatic brain injury (TBI) is a leading cause of death and disability worldwide. More than 90% of the burden is shared by low - and middle-income countries (1). Decompressive craniotomy (DC) is a surgical method performed to immediately reduce intracranial pressure (ICP) (2). DC is a known risk factor for the development of posttraumatic hydrocephalus (PTH) in the patients with TBI (3). PTH affects 11.9% - 36% of patients undergoing DC (4), is an important cause of morbidity after TBI and affects the patients' long-term outcomes (5). PTH has been recognized since Dandy's report in 1914 (6). The cause of hydrocephalus development after DC is considered to be multi-factorial, and risk factors associated with PTH still remain un-

clear (7). The treatment of PTH in patients who undergo DC is a difficult problem to solve. Shunt operation with or without cranioplasty may be done simultaneously aiming to correct the underlying problems. Therefore, it is important to prevent the PTH for the good outcomes of the patients with DC (8). The purpose of this study was to evaluate the development of PTH requiring ventriculoperitoneal (VP) shunt after DC for TBI.

2. MATERIALS AND METHODS**Search strategies**

Four databases (PubMed, Web of Science, Scopus, and Cochrane Library) were searched using key terms "hydrocephalus" and "decompressive craniectomy" and ("head trauma" or "traumatic head" or "head inju-

© 2018 Reza Fattahian, Seyed Reza Bagheri, Masoud Sadeghi

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

The first author, publication year	Country	Study Design	No. of patients undergoing DC	No. of patients with Posttraumatic Hydrocephalus	Mean initial (preoperative or admission) Glasgow Coma score	Mean age/range of patients undergoing DC, year	%male in patients undergoing DC
Yang, 2003 (18)]	China	Retrospective	68	20	-	-	-
Aarabi, 2006 (10)	USA	Retrospective cohort	50	5	3-15	25.3/>18	66
Kan, 2006 (11)	USA	Retrospective cohort	51	20	4.6	6.6/<18	64.7
Jagannathan, 2007 (12)	USA	Retrospective	17	5	4.6	11.9/2-19	65
Choi, 2008 (8)	Korea	Retrospective	55	13	9.2	49/>30	-
Morgalla, 2008 (21)	Germany	Retrospective	33	4	≤8	36.3/13-60	60.6
Yang, 2008 (19)	China	Retrospective	108	10	Range: 3 to >9	44.3/<18 to >60	68.5
Aarabi, 2009 (13)	USA	Retrospective cohort	54	7	6.6	Median: 32 /16-68	83.3
Ban, 2010 (16)	Korea	Retrospective	89	10	7.2	51.4/4-82	88.8
Honeybul, 2010 (23)	Australia	Retrospective cohort	41	4	≤8	32.1/-	83
Honeybul, 2011 (24)	Australia	Retrospective	164	23	-	-	-
Malmivaara, 2011 (25)	Finland	Retrospective cohort	54	6	≤8:83% > 8:17%	37 years/13-65	81.5
Su, 2011(26)	Taiwan	Retrospective	149	35	≤8	-	-
De Bonis, 2013 (3)	Italy	Retrospective cohort	64	19	7.3	37.9/16-80	79.7
Saade, 2014 (27)	Brazil	Retrospective	56	6	4/5 critical:51.7%& >5: 27: 48.3%	-/18-65	83.9
Sedney, 2014 (14)	USA	Retrospective	20	2	3.8	37.8/≥ 18	90
Zeiler, 2014 (28)	Canada	Retrospective	20	4	6.5	44.1/19-72	75
Grille, 2015 (29)	Uruguay	Retrospective	64	4	7	31/-	79
Ki, 2015(17)	Korea	Retrospective	92	24	Range: 3 to >8	52.8/-	74
Pechmann, 2015 (20)	Germany	Retrospective cohort	12	5	4.5	8.5/2-14	67
Sinha, 2015 (30)	India	Retrospective cohort	944	69	≤8	Median:32/-	82.7
Yuan, 2015 (20)	China	Retrospective cohort	62	16	-	50/>18	-
Kinoshita, 2016 (31)	Japan	Retrospective cohort	39	6	7	-/≥60	71.8
Jehan, 2017 (15)	USA	Retrospective cohort	33	3	Median: 9	48.8/>18	69.7
Kim, 2017 (5)	Korea	Retrospective	63	34	-	53.7/7-85	77.8

Table 1: The characteristics of studies included in meta-analysis (n=25). Abbreviation: DC; decompressive craniectomy

ry”or“brain injury”or“brain trauma”or“traumatic brain”). The databases were searched from 1983 to April 2018.

Study selection and selection criteria

The studies on the prevalence of PTH requiring VP shunt after DC in patients with TBI were selected without language restriction. The studies were selected for analysis if they a) were retrospective studies; b) included patients with TBI for any reason; c) showed the prevalence of PTH after DC alone; d) included patients with any age; e) diagnosed hydrocephalus by computed tomography scan, and f) included patients with any Glasgow Coma Score (GCS).

Data extraction

One author (M.S) checked all studies and selected the eligible ones. Then, the author extracted the relevant data, including the first author, publication year, country, number of patients undergoing DC, number of patients with hydrocephalus, GCS of patients, age/range of pa-

tients, and percentage of males in patients undergoing DC. Another author (R.F) re-checked the data and disagreements were resolved via conversation by both. The definition of hydrocephalus was based on the study of Honeybul and Ho (9).

Statistical analyses

A random-effects meta-analysis using event rate (ER) and 95% confidence intervals (CIs) was done by Review Manager 5.3 (RevMan 5.3, The Cochrane Collaboration, Oxford, United Kingdom). I^2 statistic was used to determine heterogeneity between the studies, and it was considered heterogeneous if $P<0.1$, and P -value (2-sided) less than 0.05 was considered statistically significant in other analyses. The publication bias was assessed through funnel plot analysis using Begg's and Egger's tests.




Figure 1. The flowchart of the study

3. RESULTS

Out of 355 studies retrieved from the databases, after removing the duplicate studies and excluding the irrelevant ones, the full-texts of 44 studies were assessed for eligibility. Nineteen studies were excluded with reasons (Figure 1). At last, 25 studies were included and analyzed in the meta-analysis.

The characteristics of 25 studies included in meta-analysis are shown in Table 1. Six studies (10-15) were reported from the USA, four studies (5,8,16,17) from Korea, three studies (18-20) from China, two studies (21, 22) from Germany, two studies (23, 24) from Australia, and one study from Italy (3), Finland (25), Taiwan (26), Brazil (27), Canada (28), Uruguay (29), India (30), and Japan (31). The studies included 2402 patients undergoing DC for TBI, 354 of who had PTH. Preoperative or admission GCS, mean age/range, and percentage of males in

the patients undergoing DC are shown in Table 1. In all studies, the percentage of males was higher than females.

Event rate of hydrocephalus (without age restriction)

The pooled ER of PTH in the patients undergoing DC for TBI was 17.7% [95%CI: 13.2 to 23.4%; $P<0.001$] and $I^2=86.6\%$ (Figure 2).

Event rate of hydrocephalus (adult vs. children)

The pooled analysis showed that ER of PTH was 13% in adults [95%CI: 9 to 18.5%; $P<0.001$; $I^2=74.7\%$] and 37.6% in children [95%CI: 27.79 to 48.7%; $P=0.029$; $I^2=0\%$] (Figure 3). Therefore, ER of PTH in children undergoing DC for TB was around three times more than adults.

Publication bias

Egger's and Begg's tests did not show any publication bias in each analysis (Figure 4).

4. DISCUSSION

TBI is one of the most common causes of death among young people in the industrialized countries (32). This meta-analysis showed that the ER of PTH in patients with TBI and undergoing DC was significantly high (17.7%). In addition, the ER of PTH was higher in children than adults (37.6% vs. 13%).

Among the studies included in meta-analysis, Grille et al. (29) reported the lowest rate (6.3%) and Kim et al. (5) reported the highest rate (54%). The low rate of hydrocephalus in Grille's study (29) was perhaps due to under diagnosis or utilization of different diagnostic criteria according to international references, confirmed by the study of Honeybul and Ho (24). But in the study of Kim et al., (5) among the coexisting hemorrhages at the first admission, the maximal thickness of traumatic subdural hemorrhage and the extent of midline shift were significantly larger in the group with PTH compared to the group without PTH, and another study confirmed the extent of midline shift (3). The study of Honeybul and Ho (9) found that the development of hydrocephalus was associated with low admission GCS score and subdural hygroma rather than craniotomy close to the midline. Complications following DC for TBI were found to occur at specific times, and the poor GCSscore (≤ 8) or the severity of brain injury (11, 16, 23) and the older age (≥ 65) were found to show the high incidence of hydrocephalus (16). Also, extended craniectomy and repeated operation might play a role in the development of PTH in patients receiving DC (8). It was hypothesized that hydrocephalus might be dependent on cerebrospinal fluid (CSF) dynamic alterations (8, 33), especially in the venous reabsorption phase (33).

Therefore interhemispheric subdural hygroma may represent an epiphenomenon following DC close to the midline, and that subdural hygroma can be considered an expression of some alteration of CSF dynamics that may precede the appearance of hydrocephalus (3). One study showed that in patients with severe head trauma undergoing DC, hydrocephalus was associated with a large craniectomy area and contralateral subdural hygroma (17). Leaving the bone flap out at that time appeared to help control CP, but was associated with an increased incidence of PTH and wound complications (11). One

study (9) reported that timing of cranioplasty was not significantly associated with hydrocephalus requiring a VP shunt, but maximum ICP prior to decompression had a significant correlation with hydrocephalus. Therefore, DC causes alterations of CSF and high ICP that these changes can increase the risk of postoperative hydrocephalus.

A randomized controlled trial showed that younger age was associated with shunt-dependent hydrocephalus after DC in a recent study by Vedantam et al. (4) The presence of interhemispheric subdural hygroma and younger age were associated with shunt-dependent hydrocephalus after DC in patients with severe TBI. Low et al. (34) found that younger age was associated with PTH, while others have shown a link between older age and




Figure 2. Forest plot of random-effects of event rate of posttraumatic hydrocephalus in all patients of each age undergoing decompressive craniectomy.




Figure 3. Forest plot of random-effects of event rate of posttraumatic hydrocephalus in adults and children undergoing decompressive craniectomy.




Figure 4. Funnel plot of event rate of posttraumatic hydrocephalus in (A) all studies, (B) adults, and (C) children undergoing decompressive craniectomy.

PTH (35, 36). High-energy mechanisms of injury, higher compliance of brain tissue (37), and lack of cerebral atrophy could contribute to an increased risk of symptomatic hydrocephalus in younger patients after DC (4). A prospective cohort among the patients that survived long enough showed the higher incidence of hydrocephalus requiring VP shunt in patients who had bilateral DC compared to unilateral DC (21% vs. 9%; $P>0.05$) (38). Therefore, GCS score, alterations of CSF, ICP, and age may play significant roles in the development of PTH in patients receiving DC.

Limitations

The age range, GCS, and time of follow-up, DC procedure, and type of trauma were different between the studies.

5. CONCLUSIONS

The present study demonstrated that DC after TBI was associated with the development of PTH especially in children compared to adults that several factors can impact on the incidence of PTH. Therefore, randomized controlled clinical trials are needed to confirm these conclusions by controlling these factors affecting the development of PTH.

- **Authors Contributions:** Conception, design, acquisition of data, and analysis and interpretation of data: Reza Fattahian and Masoud Sadeghi; Drafting the article: Masoud Sadeghi; Critically revising the article: Seyed Reza Bagheri.
- **Conflict of interest:** We have no conflicts of interest to disclose.

REFERENCES

1. Hofman K, Primack A, Keusch G, Hrynokow S. Addressing the growing burden of trauma and injury in low-and middle-income countries. *Am J Public Health*. 2005; 95(1): 13-17.
2. Faleiro RM, Faleiro LC, Caetano E, Gomide I, Pita C, Coelho G, et al. Decompressive craniotomy: prognostic factors and complications in 89 patients. *Arq Neuropsiquiatr*. 2008; 66(2B): 369-373.
3. De Bonis P, Sturiale CL, Anile C, Gaudino S, Mangiola A, Martucci M, et al. Decompressive craniectomy, interhemispheric hygroma and hydrocephalus: a timeline of events? *Clin Neurol Neurosurg*. 2013; 115(8): 1308-1312.
4. Vedantam A, Yamal JM, Hwang H, Robertson CS, Gopinath SP. Factors associated with shunt-dependent hydrocephalus after decompressive craniectomy for traumatic brain injury. *J Neurosurg*. 2018; 128(5): 1547-1552.
5. Kim H, Lee HS, Ahn SY, Park SC, Huh W. Factors Associated Postoperative Hydrocephalus in Patients with Traumatic Acute Subdural Hemorrhage. *J Korean Neurosurg Soc*. 2017; 60(6): 730-737.
6. Dandy W, Blackfan KD. Internal hydrocephalus. An experimental, clinical and pathological study. *Am J Dis Child*. 1914; 8(6): 406-482.
7. Wani AA, Ramzan AU, Tanki H, Malik NK, Dar BA. Hydrocephalus after decompressive craniotomy: a case series. *Pediatr Neurosurg*. 2013; 49(5): 287-291.
8. Choi I, Park HK, Chang JC, Cho SJ, Choi SK, Byun BJ. Clinical factors for the development of posttraumatic hydrocephalus after decompressive craniectomy. *J Korean Neurosurg Soc*. 2008; 43(5): 227-231.
9. Honeybul S, Ho KM. Incidence and risk factors for post-traumatic hydrocephalus following decompressive craniectomy for intractable intracranial hypertension and evacuation of mass lesions. *J Neurotrauma*. 2012; 29(10): 1872-1878.
10. Aarabi B, Hesdorffer DC, Ahn ES, Aresco C, Scalea TM, Eisenberg HM. Outcome following decompressive craniectomy for malignant swelling due to severe head injury. *J Neurosurg*. 2006; 104(4): 469-479.
11. Kan P, Amini A, Hansen K, White GL Jr, Brockmeyer DL, Walker ML, et al. Outcomes after decompressive craniectomy for severe traumatic brain injury in children. *J Neurosurg*. 2006; 105(5 Suppl): 337-342.
12. Jagannathan J, Okonkwo DO, Dumont AS, Ahmed H, Bahari A, Prevedello DM, et al. Outcome following decompressive craniectomy in children with severe traumatic brain injury: a 10-year single-center experience with long-term followup. *J Neurosurg*. 2007; 106(4 Suppl): 268-275.
13. Aarabi B, Hesdorffer DC, Simard JM, Ahn ES, Aresco C, Eisenberg HM, et al. Comparative study of decompressive craniectomy after mass lesion evacuation in severe head injury. *Neurosurgery*. 2009; 64(5): 927-939.
14. Sedney CL, Julien T, Manon J, Wilson A. The effect of craniectomy size on mortality, outcome, and complications after decompressive craniectomy at a rural trauma center. *J Neurosci Rural Pract*. 2014; 5(3): 212-217.
15. Jehan F, Azim A, Rhee P, Khan M, Gries L, O'Keeffe T, et al. Decompressive craniectomy versus craniotomy only for intracranial hemorrhage evacuation: A propensity matched study. *J Trauma Acute Care Surg*. 2017; 83(6): 1148-1153.
16. Ban SP, Son YJ, Yang HJ, Chung YS, Lee SH, Han DH. Anal-

- ysis of complications following decompressive craniectomy for traumatic brain injury. *J Korean Neurosurg Soc.* 2010; 48(3): 244-250.
17. Ki HJ, Lee HJ, Lee HJ, Yi JS, Yang JH, Lee IW. The Risk Factors for Hydrocephalus and Subdural Hygroma after Decompressive Craniectomy in Head Injured Patients. *J Korean Neurosurg Soc.* 2015; 58(3): 254-561.
 18. Yang XJ, Hong GL, Su SB, Yang SY. Complications induced by decompressive craniectomies after traumatic brain injury. *Chin J Traumatol.* 2003; 6(2): 99-103.
 19. Yang XF, Wen L, Shen F, Li G, Lou R, Liu WG, et al. Surgical complications secondary to decompressive craniectomy in patients with a head injury: a series of 108 consecutive cases. *Acta Neurochir (Wien).* 2008; 150(12): 1241-1247.
 20. Yuan Q, Wu X, Yu J, Sun Y, Li Z, Du Z, et al. Subdural hygroma following decompressive craniectomy or non-decompressive craniectomy in patients with traumatic brain injury: Clinical features and risk factors. *Brain Inj.* 2015; 29(7-8): 971-980.
 21. Morgalla MH, Will BE, Roser F, Tatagiba M. Do long-term results justify decompressive craniectomy after severe traumatic brain injury? *J Neurosurg.* 2008; 109(4): 685-690.
 22. Pechmann A, Anastasopoulos C, Korinthenberg R, van Velthoven-Wurster V, Kirschner J. Decompressive craniectomy after severe traumatic brain injury in children: complications and outcome. *Neuropediatrics.* 2015; 46(1): 5-12.
 23. Honeybul S. Complications of decompressive craniectomy for head injury. *J Clin Neurosci.* 2010; 17(4): 430-435.
 24. Honeybul S, Ho KM. Long-term complications of decompressive craniectomy for head injury. *J Neurotrauma.* 2011; 28(6): 929-935.
 25. Malmivaara K, Kivisaari R, Hernesniemi J, Siironen J. Cost-effectiveness of decompressive craniectomy in traumatic brain injuries. *Eur J Neurol.* 2011; 18(4): 656-662.
 26. Su TM, Lee TH, Huang YH, Su FW, Chen WF. Contralateral subdural effusion after decompressive craniectomy in patients with severe traumatic brain injury: clinical features and outcome. *J Trauma.* 2011; 71(4): 833-837.
 27. Saade N, Veiga JC, Cannoni LF, Haddad L, Araújo JL. Evaluation of prognostic factors of decompressive craniectomy in the treatment of severe traumatic brain injury. *Rev Col Bras Cir.* 2014; 41(4): 256-262.
 28. Zeiler FA, West M. Decompressive craniectomy: contralateral lesions and metabolic abnormalities. *Can J Neurol Sci.* 2014; 41(3): 350-356.
 29. Grille P, Tommasino N. Decompressive craniectomy in severe traumatic brain injury: prognostic factors and complications. *Rev Bras Ter Intensiva.* 2015; 27(2): 113-118.
 30. Sinha S, Raheja A, Garg M, Moorthy S, Agrawal D, Gupta DK, Satyarthee GD, Singh PK, Borkar SA, Gurjar H, Tandon V, Pandey RM, Sharma BS. Decompressive craniectomy in traumatic brain injury: A single-center, multivariate analysis of 1,236 patients at a tertiary care hospital in India. *NeuroIndia.* 2015; 63(2): 175-183.
 31. Kinoshita T, Yoshiya K, Fujimoto Y, Kajikawa R, Kiguchi T, Hara M, et al. Decompressive Craniectomy in Conjunction With Evacuation of Intracranial Hemorrhagic Lesions Is Associated With Worse Outcomes in Elderly Patients With Traumatic Brain Injury: A Propensity Score Analysis. *World Neurosurg.* 2016; 89: 187-192.
 32. Martins ET, Linhares MN, Sousa DS, Schroeder HK, Meinerz J, Rigo L A, et al. Mortality in severe traumatic brain injury : a multivariate analysis of 748 Brazilian patients from Florianópolis City. *J Trauma.* 2009; 67(1): 85-90.
 33. De Bonis P, Pompucci A, Mangiola A, Rigante L, Anile C. Post-traumatic hydrocephalus after decompressive craniectomy: an underestimated risk factor. *J Neurotrauma.* 2010; 27(11): 1965-1970.
 34. Low CY, Low YY, Lee KK, Chan SP, Ang BT. Post-traumatic hydrocephalus after ventricular shunt placement in a Singaporean neurosurgical unit. *J Clin Neurosci.* 2013; 20(6): 867-872.
 35. Jiao QF, Liu Z, Li S, Zhou LX, Li SZ, Tian W, et al. Influencing factors for posttraumatic hydrocephalus in patients suffering from severe traumatic brain injuries. *Chin J Traumatol.* 2007; 10: 159-162.
 36. Kaen A, Jimenez-Roldan L, Alday R, Gomez PA, Lagares A, Alén JF, et al. Interhemispheric hygroma after decompressive craniectomy: does it predict posttraumatic hydrocephalus? *J Neurosurg.* 2010; 113(6): 1287-1293.
 37. Kiening KL, Schoening W, Unterberg AW, Stover JF, Citerio G, Enblad P, et al. Assessment of the relationship between age and continuous intracranial compliance. *Acta Neurochir Suppl.* 2005; 95: 293-297.
 38. Honeybul S, Ho KM. Decompressive craniectomy for severe traumatic brain injury: the relationship between surgical complications and the prediction of an unfavourable outcome. *Injury.* 2014; 45(9): 1332-1339.