



RESEARCH ARTICLE

Meditation and Mental Health (Depression, Anxiety, and Stress) in Saudi Arabia

NOARA ALHUSSEINI^{1*}, SAMI ALMUSTANYIR, MD², DALIA HAMDAN³, RAGHAD HIJAZI⁴, REEMA ALDHALAAN⁵, AREEJ ABDUL GHANI ALFATTANI, MPH, CCRP⁶, DR. AAMIR OMAIR⁷

¹Associate Director MPH program, Senior Lecturer, College of Medicine, Alfaisal University, Riyadh, Saudi Arabia ,
Email: nalhusseini@alfaisal.edu / alhusseini.n@gmail.com

²College of Medicine, Alfaisal University, Riyadh, Saudi Arabia , Email: saalmustanyir@alfaisal.edu

³College of Medicine, Alfaisal University, Riyadh, Saudi Arabia , Email: dnhamdan@alfaisal.edu

⁴College of Medicine, Alfaisal University, Riyadh, Saudi Arabia , Email: ramhijazi@alfaisal.edu

⁵College of Medicine, Alfaisal University, Riyadh, Saudi Arabia , Email: raldhalaan@alfaisal.edu

⁶Research Associate, Department of Biostatistics, Epidemiology and Scientific Computing, King Faisal Specialist Hospital and Research Center, Riyadh, Saudi Arabia, Email: aralfattani@kfshrc.edu.sa

⁷Lecturer, Research Unit, Dept. of Med Education, College of Medicine, King Saud bin Abdulaziz University for Health Sciences, Riyadh, Saudi Arabia, Email: Amir.omair@gmail.com

*Corresponding Author

ABSTRACT

Introduction: Meditation encompasses any practice that connects an individual to their inner self. All these practices essentially serve to relax and divert both the mind and the body away from worries and concerns. The popularity of meditation has grown tremendously in the last century. Due to its correlation with mental health, research efforts spiked to uncover an association.

Purpose: The main objective is to measure the frequency of meditation among the Saudi population and to understand its impact on mental health (depression, anxiety, and stress).

Methods: We used a cross-sectional design among the population in Saudi Arabia. A validated online questionnaire was distributed via social media channels using convenience sampling. Chi-square tests were used to determine associations between meditation, demographics, depression, anxiety, and stress.

Results: A total of 902 respondents answered the survey. The majority (75%) reported practicing some form of meditation and the most common meditation practice was exercise, which was reported by 390 (43%). The proportion of those with severe/extremely severe depression was higher in the group that did not meditate (38%) as compared to 29% in the group that meditated ($p=0.03$). A similar difference was seen for the stress levels, with 38% of the non-meditating group having severe/extreme level of stress as compared to 28% in the meditating group ($p=0.03$). There was no significant difference in the anxiety level between the two groups ($p=0.38$).

Conclusion: Meditation practice frequency is relatively high in Saudi Arabia. Our study uncovered an association between meditation and mental health status. The proportion of those with severe/extremely severe depression and stress was higher in the group that did not meditate. Meditation must be given higher priority in the management of mental health cases, especially in the early stages. Due to the limited meditation studies among the Saudi population, further research is essential to better understand the relationship between mental health and meditation.

KEYWORDS:

Meditation, Depression, Anxiety, Stress, Mental Health Saudi Arabia

ARTICLE HISTORY:

Received June 07, 2021
Accepted July 12, 2021
Published Sep 15, 2021

DOI:

10.5455/jcmr.2021.12.02.21

VOLUME: 12

ISSUE: 2

ISSN: 2146-8397

INTRODUCTION

Meditation takes on many different forms and does not have a unified definition. The most widely agreed-upon notion is that meditation encompasses any practice that connects an individual to their inner self. All these practices essentially serve to relax and divert both the mind and the body away from worries and concerns (Sharma, 2015). The ultimate goal of meditation is to focus on the present moment and block any external distractions. Meditation is categorized into seven main groups: mindful observation, body-centered meditation, visual concentration, contemplation, affect-centered meditation, mantra meditation, and meditation with movement (Matko & Sedlmeier, 2019).

The benefits of meditation are a consequence of a neurophysiological change; meditation acts on the autonomic nervous system by reducing the activity of the sympathetic nervous system and increasing the activity of the parasympathetic nervous system. This will result in an overall decrease in the release of stress hormones like catecholamines and cortisol occur (Rajaraman, 2013).

Concentration meditation and mindfulness meditation are two essential forms of meditation. Concentration meditation intends to willfully direct the mind's attention onto a particular object only, such as one's breath, the rhythm of a chant, or an illusory object. Mindfulness meditation intends to increase one's awareness about the current mental status (Latham, 2016). Breathing is a form of meditation. Furthermore, Pranayama is a regular practice of breathing exercises. In a study that measured the effect of pranayama on the function of both the sympathetic and parasympathetic systems, it was found to cause sympathovagal balance, which served to relax the body and mind. This was found to decrease mental distress and increase well-being (Singh¹ et al., 2019).

Creating art is also considered a form of meditation, which presents a wide scope of meditation methods. A newly developed therapy known as meditation-based art therapy (MBAT) has been found to reduce stress levels. A study used functional magnetic resonance imaging (fMRI) on individuals that practiced MBAT found that cerebral blood flow decreased during stressful situations (Hinchey, 2018).

Meditation practices are linked to improving meditators' mental health status and reducing stress levels. A correlation between meditation and improved intelligence level has been explored as well. In addition, a study that aimed to identify the immediate and long-term effects found an increase in both IQ and cognitive abilities (Prakash et al., 2012).

Purpose and Significance

The objective is to measure the frequency of meditation among the Saudi population and to understand its impact on mental health (depression, anxiety, and stress). There are limited peer-reviewed studies about meditation and mental health among the Saudi population. This study will be novel as it will add value to the current literature.

METHODS

A cross-sectional design was used among Saudi Arabia's residents, including adults above 18 years old, Saudis, and non-Saudis. An online questionnaire was created using Google docs and distributed via social media channels such as WhatsApp, LinkedIn, Facebook, Twitter, and Snapchat using convenience sampling. The estimated annual population of Saudi Arabia in 2020 was around 34.8 million. Our sample size was determined to be 384 at the 95% confidence and a 5% margin of error, with an expected outcome response of 50%. We aimed for higher response to ensure generalizability.

A validated questionnaire for the meditation (Soler et al., 2014) and the Depression Anxiety Stress Scales questionnaire (DASS) (Australia, 2018) were both used and adjusted to fit the objective of our study. The questionnaire included three main sections. The first section included demographic questions, including age, gender, nationality, employment status, marital status, and monthly income. The second section included questions related to meditation practices including, type and frequency of meditation. The last section covered (DASS). The questions were translated word by word to Arabic by a translator. It was distributed in English and Arabic. Face and content validity were achieved by experts in the field and a translator. The questionnaire was anonymous to ensure the respondents' privacy and confidentiality. Access to the surveys' responses was limited to the investigators. The Institutional Review Board (IRB) at Alfaisal University approved this study.

The authors used SPSS version 24 to analyze the data. The authors presented descriptive statistics as frequencies and percentages for the categorical variables. The authors used Chi-square test to determine an association between meditation practice and demographics, as well as with the depression, anxiety, and stress categories. The significance level for the statistical tests was set at 0.05.

RESULTS

The online survey included a total of 902 respondents. The majority of these were females (74%), aged between 18 to 29 years (59%), and Saudi nationals (84%), as shown in Table 1. The majority of them were single (65%) and not employed (59%); of those who reported their monthly income, 283 (32%) did not have a monthly income, and 202 (23%) had a monthly income of less than SR 10,000 (US\$ 2650).

There were a total of 674 (75%) respondents who reported practicing some form of meditation; of these, 641 respondents reported their frequency of meditation. Almost half of them, i.e., 279 (44%), practiced some meditation on a daily basis, while another 232 (36%) reported doing some meditation weekly. The majority of them, i.e., 385 (60%), practiced meditation for up to <30 minutes per session, and most (64%) of them were doing meditation for more than six months (Fig 1). The most common meditation practice was exercise, which was reported by 390 (43%) of the 902 respondents, followed by 316 (35%) practicing breathing, 254 (28%) for faith-based

practices, and 237 (26%) for yoga (Fig 2).

Table 2 shows the comparison of demographic characteristics between those who practiced meditation (n=674) and those who did not meditate (n=219). Of those who did not meditate, 146 (67%) were aged 18-29 years as compared to 385 (57%) in the group that meditated ($p=0.04$). There was a greater proportion of females in the group that practiced meditation (76%), while there were 68% females in the group that did not meditate ($p=0.03$). There were no other significant differences for nationality, employment status, or monthly income between those who practiced meditation and those who did not ($p>0.05$).

Table 3 shows the comparison of depression, anxiety, and stress levels between those who meditate (n=647) and those who do not meditate (n=191). It was found that the proportion of those with severe/extremely severe depression was higher in the group that did not meditate (38%) as compared to 29% in the group that meditated ($p=0.03$). A similar difference was seen for the stress levels, with 38% of the not meditating group having severe/extreme levels of stress as compared to 28% in the meditating group ($p=0.03$). There was no significant difference in the anxiety level between the two groups ($p=0.38$).

DISCUSSION

The popularity of meditation has grown tremendously in the last century. Due to its correlation with mental health, research efforts spiked to uncover an association (Kang et al., 2014; Kok & Singer, 2017; Sharma, 2015). Therefore, the main objective of this study is to measure the frequency of meditation among the Saudi population and to understand its impact on mental health (depression, anxiety, and stress).

The prevalence of meditation is high in Saudi Arabia; our findings indicated that 674 (75%) of respondents reported practicing some form of meditation. Similarly, others found unconventional treatment methods (such as meditation) to be more accepted by Saudis; 52% of 2114 participants reported practicing complementary and alternative medicine (CAM) (AlShehri et al., 2020). A study used the National Health Interview Survey NHIS of USA in 2012, including 34,525 individuals; it concluded that the prevalence of meditation was 5.2% lifetime and 4.1% in a 12 months period (Cramer, Hall, et al., 2016). The prevalence of meditation practice increased globally as more individuals are discovering its value. It was estimated that almost 21 million USA adults practiced yoga in the last 12 months (Cramer, Ward, et al., 2016).

Our results showed that 279 (44%) of the respondents practiced some form of meditation daily, while another 232 (36%) reported doing some meditation on a weekly basis.

The majority of them, i.e., 385 (60%), practiced meditation for up to <30 minutes per session, and most (64%) of them were practicing meditation for more than six months. Lemay et al., (2019) found that mindful meditation and yoga activities as infrequent as thirty minutes once a week have been sufficient to significantly reduce stress and anxiety levels (Lemay et al.,

2019).

Since meditation is practiced in many forms, we found that the most common meditation practice was exercise, which was reported by 390 (43%) of the 902 respondents, 315 (35%) practiced breathing, 254 (28%) used faith-based or spiritual practices, followed by 237 (26%) for yoga. A study in the United States found spiritual meditation to have a 3.1% prevalence, mindfulness meditation to have a 1.9% prevalence, and mantra meditation to have a 1.6% prevalence (Burke et al., 2017). Another study found an incline from 8.9% (n = 3,080) in 2012 to 13.3 % (n = 3,552) in 2017 in the prevalence of yoga and breathing meditation (Yobam) amongst study participants (Schröter & Cramer, 2021). The differences in meditation forms practiced could be due to the differences in the cultures.

Our study indicated that for those who did not meditate, 146 (67%) were aged 18-29 years as compared to 385 (57%) in the group that meditated ($p=0.04$). It was seen that there was a greater proportion of females in the group that practiced meditation (76%), while there were 68% females in the group that did not meditate ($p=0.03$). Our findings are in concordance with other studies that found that there is a higher prevalence of women using complementary and alternative (CAM) methods such as yoga and tai chi. They also found that women perceived it to be more helpful in improving their overall sense of well-being than their male counterparts (Upchurch & Johnson, 2019). Others found that those who meditate generally adopt health consciousness and physically active lifestyles when compared to non-meditators. Higher utilization of meditation was found amongst females with some forms of functional limitations or health complaints such as lower back pain and depression in the past 12 months (Burke et al., 2017). Some studies found that individuals meditated to improve their general wellbeing (76.2%), to improve their energy (60.0%), or to help them focus (50.0%). The health problems for which people meditated were anxiety (29.2%), stress (21.6%), and depression (17.8%) (Cramer, Hall, et al., 2016). A study found that both genders meditated to manage stress. More commonalities found amongst the genders were that both are more likely to be unmarried, have earned a college level of education or higher, and live with some form of mental distress (Upchurch & Johnson, 2019).

We found that the proportion of those with severe/extremely severe depression was higher in the group that did not meditate (38%) as compared to 29% in the group that meditated ($p=0.03$). A randomized controlled trial aimed to explore the effect of mindfulness meditation on patients with a lifetime history of depression showed a significant reduction in depressive symptoms (Winnebeck et al., 2017). A meta-analysis examined the effects of mindfulness meditation interventions (MMI) on depression among older adults. It included 1,076 participants and concluded that MMI significantly improved depressive symptoms (Reangsing et al., 2020). Studies confirmed that mindfulness meditation could be used as an alternative treatment for the early stages of depression (Kyselova et al., 2018). Similarly, it was found that

meditation has a significant effect in reducing depression, especially amongst university students (González-Valero et al., 2019). However, another study observed the impact of Yoga Nidra and meditation on anxiety and depression among 60 college professors indicated better anxiety results for the intervention groups with no significant improvement for the Beck Depression Inventory (Ferreira-Vorkapic et al., 2018), which can be due to the complexity of depression or its late stages. A study indicated that exercise-based yoga was not effectively reducing the symptoms of depression; however, integrative styles that emphasize meditation were found to be effective (Gong et al., 2015).

Our findings indicated a difference in the stress levels, with 38% of the non-meditating group having severe/extreme levels of stress as compared to 28% in the meditating group ($p=0.03$). A randomized control study confirms our findings as it shows that Internet Mindfulness Meditation Intervention (IMMI) improved depression, perceived stress, insomnia and pain severity (Wahbeh, 2018). A meta-analysis of 34 studies revealed that meditation practices had the largest impact on the reduction of stress levels. A moderately positive impact uncovered a decrease in anxiety symptoms and a small but significant impact on depressive symptoms. Mindful meditation, in particular, even for short durations as five to eight weeks, has been found to show the greatest impact on mental health (González-Valero et al., 2019). Moreover, evidence of improved memory, quality of sleep, and stress management have also been reported among meditators compared to non-meditators (Burke et al., 2017).

No significant difference in the anxiety level between the two groups ($p=0.38$) was found. Saeed et al. (2019) found that nonconventional interventions, including meditation, can improve depression and anxiety disorders; however, positive findings are less common in individuals with anxiety disorders (Saeed et al., 2019). Another study on a group of university students in their early twenties found that women reported significant decreases in negative affect after participating in mindful meditation for a 12-week period. Men, on the other hand, did not have significant improvements in self-reported mood despite there being no differences in hours meditated and matching of baseline effect at the beginning of the study (Rojiani et al., 2017). Not all studies are in favor of mindful meditation as a solution to anxiety syndromes. A study investigating the efficacy of mindful meditation on hemodialysis patients found that despite up to 50% of patients experienced depression and anxiety, no significant effects were found from the implementation of meditation by these patients. This intervention was reported to be enjoyable by the patients but fail in terms of long-term efficacy in maintaining a significant drop in anxiety levels when assessed by questionnaires. These findings may suggest that patients with milder forms of anxiety would be more eligible candidates for mindful meditation treatment for anxiety (Thomas et al., 2017). Some studies have even reported instances in which episodes of psychosis, panic, anxiety, and traumatic memory re-experiencing were induced by meditation sessions. These studies claim that individuals with serious illnesses might be

misled by the over-exaggeration of the true efficacy of mindful meditation and that more traditional methods such as pharmacotherapy should not be replaced by meditation practices (Van Dam et al., 2018). Many studies have shown positive views of meditation in treating depression, however, there is a body of research that contradicts such claims. A systematic review of 83 articles that analyzed adverse meditation events (MAE) noted that the most common adverse effects were psychiatric MAE, which was found in 40 articles (49%). The most common symptoms were anxiety (18 studies) and depression (15 studies). Such adverse effects were found even in those with no previous psychiatric disorders. This emphasizes the importance of considering the negative impact of meditation (Farias et al., 2020). Similarly, a review of observational and experimental studies aimed to explore the negative impact of meditation. Upon reviewing 39 articles, among the most common adverse effects noted were: emotional difficulties, a decline in memory, delusions, and distorted sense of self. This further emphasizes the potential harm that meditation practice can pose (Lambert et al., 2021).

CONCLUSION

Meditation practice frequency is relatively high in Saudi Arabia. Our study uncovered an association between meditation and mental health status. The proportion of those with severe/extremely severe depression and stress was higher in the group that did not meditate. There was no significant difference in the anxiety level between the two groups. Furthermore, meditation must be given higher priority in the management of mental health problems, especially in the early stages. Due to the cross-sectional nature of the study, causal inference cannot be obtained. Using DASS as a measure of depression, stress, and anxiety does not indicate a clinical diagnosis. Due to the limited meditation studies among the Saudi population, further research is essential to better understand the relationship between mental health and meditation.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

REFERENCES

1. AlShehri, S. D., AbdulHameed, R. M., Taha, A. Z., Almusalmi, A. M., Almulaify, M. S., & Alkhabbaz, F. L. (2020, May-Aug). Complementary and alternative medicine practice and perceptions of attendees of primary care centers in Eastern Saudi Arabia. *Journal of family & community medicine*, 27(2), 120-124. https://doi.org/10.4103/jfcm.JFCM_218_19
2. Australia, P. F. o. (2018). DASS. <http://www2.psy.unsw.edu.au/dass/>
3. Burke, A., Lam, C. N., Stussman, B., & Yang, H. (2017). Prevalence and patterns of use of mantra, mindfulness and spiritual meditation among adults in the United States. *BMC Complementary and Alternative Medicine*, 17(1), 1-18.
4. Cramer, H., Hall, H., Leach, M., Frawley, J., Zhang, Y., Leung, B., Adams, J., & Lauche, R. (2016). Prevalence, patterns, and predictors of meditation use among US adults: A nationally representative survey. *Scientific reports*, 6(1), 1-9.
5. Cramer, H., Ward, L., Steel, A., Lauche, R., Dobos, G., & Zhang,

- Y. (2016). Prevalence, patterns, and predictors of yoga use: results of a US nationally representative survey. *American journal of preventive medicine*, 50(2), 230-235.
6. Farias, M., Maraldi, E., Wallenkampf, K., & Lucchetti, G. (2020). Adverse events in meditation practices and meditation-based therapies: a systematic review. *Acta Psychiatrica Scandinavica*, 142(5), 374-393.
 7. Ferreira-Vorkapic, C., Borba-Pinheiro, C. J., Marchioro, M., & Santana, D. (2018, Sep-Dec). The Impact of Yoga Nidra and Seated Meditation on the Mental Health of College Professors. *International journal of yoga*, 11(3), 215-223. https://doi.org/10.4103/ijoy.IJOY_57_17
 8. Gong, H., Ni, C., Shen, X., Wu, T., & Jiang, C. (2015). Yoga for prenatal depression: a systematic review and meta-analysis. *BMC psychiatry*, 15(1), 1-8.
 9. González-Valero, G., Zurita-Ortega, F., Ubago-Jiménez, J. L., & Puertas-Molero, P. (2019). Use of meditation and cognitive behavioral therapies for the treatment of stress, depression and anxiety in students. A systematic review and meta-analysis. *International journal of environmental research and public health*, 16(22), 4394.
 10. Hinchey, L. M. (2018). Mindfulness-based art therapy: a review of the literature. *Inquiries Journal*, 10(05).
 11. Kang, Y., Gray, J. R., & Dovidio, J. F. (2014). The nondiscriminating heart: lovingkindness meditation training decreases implicit intergroup bias. *Journal of Experimental Psychology: General*, 143(3), 1306.
 12. Kok, B. E., & Singer, T. (2017). Effects of contemplative dyads on engagement and perceived social connectedness over 9 months of mental training: A randomized clinical trial. *Jama psychiatry*, 74(2), 126-134.
 13. Kyselova, A., Kravtsova, E., Mishchenko, D., & Chernishova, E. (2018). The Relationship between Mindfulness Meditation and Depression. *Український журнал медицини, біології та спорту*, 4(326), 304.
 14. Lambert, D., van den Berg, N., & Mendrek, A. (2021). Adverse effects of meditation: A review of observational, experimental and case studies. *Current Psychology*, 1-14.
 15. Latham, N. (2016). Meditation and self-control. *Philosophical Studies*, 173(7), 1779-1798.
 16. Lemay, V., Hoolahan, J., & Buchanan, A. (2019). Impact of a yoga and meditation intervention on students' stress and anxiety levels. *American journal of pharmaceutical education*, 83(5).
 17. Matko, K., & Sedlmeier, P. (2019). What is meditation? Proposing an empirically derived classification system. *Frontiers in psychology*, 10, 2276.
 18. Prakash, R., Rastogi, P., Dubey, I., Abhishek, P., Chaudhury, S., & Small, B. J. (2012). Long-term concentrative meditation and cognitive performance among older adults. *Aging, Neuropsychology, and Cognition*, 19(4), 479-494.
 19. Rajaraman, S. (2013). Meditation research: a comprehensive review. *International Journal of Engineering Research and Application*.
 20. Reangsing, C., Rittiwong, T., & Schneider, J. K. (2020). Effects of mindfulness meditation interventions on depression in older adults: A meta-analysis. *Aging & Mental Health*, 1-10. <https://doi.org/10.1080/13607863.2020.1793901>
 21. Rojiani, R., Santoyo, J. F., Rahrig, H., Roth, H. D., & Britton, W. B. (2017). Women benefit more than men in response to college-based meditation training. *Frontiers in psychology*, 8, 551.
 22. Saeed, S. A., Cunningham, K., & Bloch, R. M. (2019, May 15). Depression and Anxiety Disorders: Benefits of Exercise, Yoga, and Meditation. *Am Fam Physician*, 99(10), 620-627.
 23. Schröter, M., & Cramer, H. (2021). Prevalence and predictors of yogic breathing and meditation use-A nationally representative survey of US adult yoga practitioners. *Complementary Therapies in Medicine*, 56, 102617.
 24. Sharma, H. (2015). Meditation: Process and effects. *Ayu*, 36(3), 233.
 25. Singh¹, K., Garg, R., Tripathi, Y., Mann, N., & Sharma, V. (2019). Effect of Pranayama and "OM" meditation on visual reaction time, hand grip dynamometer endurance and stress in medical students. *Journal Of Medical Science And Clinical Research*, 7(7).
 26. Soler, J., Cebolla, A., Feliu-Soler, A., Demarzo, M. M., Pascual, J. C., Baños, R., & García-Campayo, J. (2014). Relationship between meditative practice and self-reported mindfulness: the MINDSENS composite index. *PloS one*, 9(1), e86622.
 27. Thomas, Z., Novak, M., Platas, S. G. T., Gautier, M., Holgin, A. P., Fox, R., Segal, M., Loooper, K. J., Lipman, M., & Selchen, S. (2017). Brief mindfulness meditation for depression and anxiety symptoms in patients undergoing hemodialysis: A pilot feasibility study. *Clinical Journal of the American Society of Nephrology*, 12(12), 2008-2015.
 28. Upchurch, D. M., & Johnson, P. J. (2019). Gender differences in prevalence, patterns, purposes, and perceived benefits of meditation practices in the United States. *Journal of Women's Health*, 28(2), 135-142.
 29. Van Dam, N. T., van Vugt, M. K., Vago, D. R., Schmalzl, L., Saron, C. D., Olendzki, A., Meissner, T., Lazar, S. W., Kerr, C. E., & Gorchov, J. (2018). Mind the hype: A critical evaluation and prescriptive agenda for research on mindfulness and meditation. *Perspectives on psychological science*, 13(1), 36-61.
 30. Wahbeh, H. (2018). Internet Mindfulness Meditation Intervention (IMMI) Improves Depression Symptoms in Older Adults. *Medicines*, 5(4), 119. <https://www.mdpi.com/2305-6320/5/4/119>
 31. Winnebeck, E., Fissler, M., Gärtner, M., Chadwick, P., & Barnhofer, T. (2017). Brief training in mindfulness meditation reduces symptoms in patients with a chronic or recurrent lifetime history of depression: A randomized controlled study. *Behaviour research and therapy*, 99, 124-130.

Table 1: Demographic characteristics of the respondents (N=902)^a

	n	(%)
Age		
18 - 29	535	59.4
30 - 44	232	25.8
> 45	133	14.8
Gender		
Female	667	74.2
Male	232	25.8
Nationality		
Saudi	755	84.3
Non - Saudi	141	15.7
Marital status		
Single	581	64.6
Married	266	29.6
Divorced / Widowed	53	5.9
Employment status		
Yes	365	40.6
No	533	59.4
Monthly income		
≤ 9,999 SR	202	22.6
10,000-19,999 SR	160	17.9
≥ 20,000 SR	91	10.2
I do not have a monthly income	283	31.7
I prefer not to answer	156	17.5

^aSome missing data for different variables - so valid percent reported

Table 2: Comparison of meditation practice by demographic characteristics

	Meditate (n=674)^a	Do Not Meditate (n=219)^a	p-value
Age (years)			
18 - 29	385 (57%)	146 (67%)	0.04*
30 - 44	186 (28%)	45 (20%)	
> 45	102 (15%)	28 (13%)	
Gender			
Female	511 (76%)	148 (68%)	0.03*
Male	163 (24%)	69 (32%)	
Nationality			
Saudi	570 (85%)	180 (83%)	0.48
Non - Saudi	101 (15%)	37 (17%)	
Marital status			
Married	203 (30%)	61 (28%)	0.03*
Single	425 (63%)	152 (70%)	
Divorced / Widowed	46 (7%)	5 (2%)	
Employment status			
Yes	281 (42%)	84 (38%)	0.39
No	391 (58%)	134 (62%)	
Monthly income			
≤ 9,999 SR	141 (21%)	58 (27%)	0.21
10,000-19,999 SR	129 (19%)	31 (14%)	
≥ 20,000 SR	72 (11%)	19 (9%)	
Doesn't have a monthly income	209 (31%)	73 (34%)	
Prefer not to answer	118 (18%)	34 (16%)	

^a Some missing data for different variables - so valid percent reported


* Significant at p<0.05

Table 3: Comparison of Depression, Anxiety, and Stress scores by meditation practice

	Meditate (n=647) ^a	Don't Meditate (n=191) ^a	p-value
Depression score			0.03*
Normal-Mild	333 (52%)	80 (43%)	
Moderate	125 (19%)	36 (19%)	
Severe-Ext Severe	185 (29%)	72 (38%)	
Anxiety score			0.38
Normal-Mild	276 (43%)	73 (38%)	
Moderate	119 (18%)	33 (17%)	
Severe-Ext Severe	252 (39%)	85 (45%)	
Stress score			0.03*
Normal-Mild	365 (56%)	93 (49%)	
Moderate	103 (16%)	26 (13%)	
Severe-Ext Severe	178 (28%)	72 (38%)	

^a Some missing data for different variables - so valid percent reported

* Significant at $p < 0.05$

**Fig.1:** Frequency and practice of meditation among the respondents (N=641)




Fig.2: Types of meditation practiced by the respondents (N=902)